

(19)



Europäisches Patentamt

European Patent Office

Office européen des brevets



(11)

**EP 0 752 248 A1**

(12)

**EUROPEAN PATENT APPLICATION**

(43) Date of publication:

**08.01.1997 Bulletin 1997/02**

(21) Application number: **96200772.0**

(22) Date of filing: **12.11.1993**

(51) Int. Cl.<sup>6</sup>: **A61K 39/395**, A61K 51/10,

C12N 15/02, C12P 21/08,

C07K 16/46

// C07K16/28

(84) Designated Contracting States:

**AT BE CH DE DK ES FR GB GR IE IT LI LU MC NL  
PT SE**

(30) Priority: **13.11.1992 US 978891**

**03.11.1993 US 149099**

(62) Application number of the earlier application in  
accordance with Art. 76 EPC: **94901444.3**

(71) Applicant: **IDEC PHARMACEUTICALS  
CORPORATION**

**San Diego, CA 92121-1104 (US)**

(72) Inventors:

- **Anderson, Darrell R.**  
**Escondido, California 92029 (US)**
- **Hanna, Nabil**  
**Olivenhain, California 92024 (US)**

• **Leonard, John E.**

**Encinitas, California 92024 (US)**

• **Newman, Roland A.**

**San Diego, California 92122 (US)**

• **Reff, Mitchell E.**

**San Diego, California 92122 (US)**

• **Rastetter, William H.**

**Rancho Santa Fe, California 92067 (US)**

(74) Representative: **Daniels, Jeffrey Nicholas et al**  
**Page White & Farrer**  
**54 Doughty Street**  
**London WC1N 2LS (GB)**

Remarks:

This application was filed on 22 - 03 - 1996 as a  
divisional application to the application mentioned  
under INID code 62.

(54) **Therapeutic application of chimeric and radiolabeled antibodies to human B lymphocyte  
restricted differentiation antigen for treatment of B cell lymphoma**

(57) Disclosed herein are therapeutic treatment pro-  
tocols designed for the treatment of B cell lymphoma.  
These protocols are based upon therapeutic strategies  
which include the use of administration of immunologi-  
cally active mouse/human chimeric anti-CD20 antibod-  
ies, radiolabeled anti-CD20 antibodies, and cooperative  
strategies comprising the use of chimeric anti-CD20  
antibodies and radiolabeled anti-CD20 antibodies.

**EP 0 752 248 A1**

**Description**

**37 C.F.R. §1.74(d)/(e) Copyright Notice**

5 A portion of the disclosure of this patent document contains material which is subject to copyright protection. The copyright owner does not object to the reproduction by anyone of the patent document or the patent disclosure, as it appears in the Patent and Trademark Office patent files or records, but otherwise reserves all copyright rights whatsoever.

10 **RELATED APPLICATIONS**

This is a Continuation-in-Part of United States Serial No. 07/978,891, filed November 13, 1992, pending. This patent document is related to United States Serial No. 07/977,691, entitled "IMPAIRED DOMINANT SELECTABLE MARKER SEQUENCE FOR ENHANCEMENT OF EXPRESSION OF CO-LINKED GENE PRODUCT AND EXPRESSION VECTOR SYSTEMS COMPRISING SAME" having U.S. Serial No. 07/977,691 (pending; filed November 13, 1992) and "IMPAIRED DOMINANT SELECTABLE MARKER SEQUENCE AND INTRONIC INSERTION STRATEGIES FOR ENHANCEMENT OF EXPRESSION OF GENE PRODUCT AND EXPRESSION VECTOR SYSTEMS COMPRISING SAME," U.S. Serial No. \_\_\_\_\_ (filed simultaneously herewith). The related patent documents are incorporated herein by reference.

TABLE OF CONTENTS

5	A.	FIELD OF THE INVENTION
	B.	BACKGROUND OF THE INVENTION
	C.	SUMMARY OF THE INVENTION
10	D.	BRIEF DESCRIPTION OF THE DRAWINGS
	E.	DETAILED DESCRIPTION OF PREFERRED EMBODIMENTS
	F.	EXAMPLES
15	I.	RADIOLABELED ANTI-CD20 ANTIBODY 2B8
	A.	Anti-CD20 Monoclonal Antibody (Murine) Production ("2B8")
	B.	Preparation of 2B8-MX-DTPA Conjugate
20	i.	MX-DTPA
	ii.	Preparation of 2B8
	iii.	Conjugation of 2B8 with MX-DTPA
25	iv.	Determination of MX-DTPA Incorporation
	v.	Immunoreactivity of 2B8-MX-DTPA
30	vi.	Preparation of Indium-[111]-labeled 2B8-MX-DTPA ("I2B8")
	vii.	Preparation of Yttrium-[90]-labeled 2B8-MX-DTPA ("Y2B8")
	C.	Non-Human Animal Studies
35	i.	Distribution of Radiolabeled 2B8-MX-DTPA
	ii.	Tumor Localization of I2B8
40	iii.	Biodistribution and Tumor Localization Studies with Radiolabeled 2B8-MX-DTPA
	D.	Human Studies
45	i.	2B8 and 2B8-MX-DTPA: Immunohistology Studies with Human Tissues
	ii.	Clinical Analysis of I2B8 (Imaging) and Y2B8 (Therapy)
50	a.	Phase I/II Clinical Trial: Single Dose Therapy Study
	b.	Phase I/II Clinical Trial: Multiple Dose Therapy Study

## II. CHIMERIC ANTI-CD20 PRODUCTION

- A. Construction of Chimeric Anti-CD20 Immunoglobulin DNA Expression Vectors
- B. Creation of Chimeric Anti-CD20 Producing CHO and SP2/0 Transfectomas
- C. Determination of Immunological Activity of Chimeric Anti-CD20 Antibodies
  - i. Human Clq Analysis
  - ii. Complement Dependent Cell Lyses
  - iii. Antibody Dependent Cellular Cytotoxicity Effector Assay

## III. DEPLETION OF B CELLS IN VIVO USING CHIMERIC ANTI-CD20

- A. Non-Human Primate Study
- B. Clinical Analysis of C2B8
  - i. Phase I/II Clinical Trial of C2B8: Single Dose Therapy Study
  - ii. Phase I/II Clinical Trial of C2B8: Multiple Dose Therapy Study

## IV. COMBINATION THERAPY: C2B8 AND Y2B8

- A. Preparation of Y2B8
- B. Preparation of C2B8
- C. Results

## V. ALTERNATIVE THERAPY STRATEGIES

## VI. DEPOSIT INFORMATION

- G. SEQUENCE LISTING
- H. CLAIMS

### **A. FIELD OF THE INVENTION**

The references to be discussed throughout this document are set forth merely for the information described therein prior to the filing dates of this document, and nothing herein is to be construed as an admission, either express or implied, that the references are "prior art" or that the inventors are not entitled to antedate such descriptions by virtue of prior inventions or priority based on earlier filed applications.

The present invention is directed to the treatment of B cell lymphoma using chimeric and radiolabeled antibodies to the B cell surface antigen Bp35 ("CD20").

### **B. BACKGROUND OF THE INVENTION**

The immune system of vertebrates (for example, primates, which include humans, apes, monkeys, etc.) consists of a number of organs and cell types which have evolved to: accurately and specifically recognize foreign microorganisms

("antigen") which invade the vertebrate-host; specifically bind to such foreign microorganisms; and, eliminate/destroy such foreign microorganisms. Lymphocytes, amongst others, are critical to the immune system. Lymphocytes are produced in the thymus, spleen and bone marrow (adult) and represent about 30% of the total white blood cells present in the circulatory system of humans (adult). There are two major sub-populations of lymphocytes: T cells and B cells. T cells are responsible for cell mediated immunity, while B cells are responsible for antibody production (humoral immunity). However, T cells and B cells can be considered as interdependent--in a typical immune response, T cells are activated when the T cell receptor binds to fragments of an antigen that are bound to major histocompatibility complex ("MHC") glycoproteins on the surface of an antigen presenting cell; such activation causes release of biological mediators ("interleukins") which, in essence, stimulate B cells to differentiate and produce antibody ("immunoglobulins") against the antigen.

Each B cell within the host expresses a different antibody on its surface - thus, one B cell will express antibody specific for one antigen, while another B cell will express antibody specific for a different antigen. Accordingly, B cells are quite diverse, and this diversity is critical to the immune system. In humans, each B cell can produce an enormous number of antibody molecules (*ie* about  $10^7$  to  $10^8$ ). Such antibody production most typically ceases (or substantially decreases) when the foreign antigen has been neutralized. Occasionally, however, proliferation of a particular B cell will continue unabated; such proliferation can result in a cancer referred to as "B cell lymphoma."

T cells and B cells both comprise cell surface proteins which can be utilized as "markers" for differentiation and identification. One such human B cell marker is the human B lymphocyte-restricted differentiation antigen Bp35, referred to as "CD20." CD20 is expressed during early pre-B cell development and remains until plasma cell differentiation. Specifically, the CD20 molecule may regulate a step in the activation process which is required for cell cycle initiation and differentiation and is usually expressed at very high levels on neoplastic ("tumor") B cells. CD20, by definition, is present on both "normal" B cells as well as "malignant" B cells, *ie* those B cells whose unabated proliferation can lead to B cell lymphoma. Thus, the CD20 surface antigen has the potential of serving as a candidate for "targeting" of B cell lymphomas.

In essence, such targeting can be generalized as follows: antihodies specific to the CD20 surface antigen of B cells are, *eg* injected into a patient. These anti-CD20 antibodies specifically bind to the CD20 cell surface antigen of (ostensibly) both normal and malignant B cells; the anti-CD20 antibody bound to the CD20 surface antigen may lead to the destruction and depletion of neoplastic B cells. Additionally, chemical agents or radioactive labels having the potential to destroy the tumor can be conjugated to the anti-CD20 antibody such that the agent is specifically "delivered" to, *e.g.* the neoplastic B cells. Irrespective of the approach, a primary goal is to destroy the tumor: the specific approach can be determined by the particular anti-CD20 antibody which is utilized and, thus, the available approaches to targeting the CD20 antigen can vary considerably.

For example, attempts at such targeting of CD20 surface antigen have been reported. Murine (mouse) monoclonal antibody 1F5 (an anti-CD20 antibody) was reportedly administered by continuous intravenous infusion to B cell lymphoma patients. Extremely high levels (>2 grams) of 1F5 were reportedly required to deplete circulating tumor cells, and the results were described as being "transient." Press *et al.*, "Monoclonal Antibody 1F5 (Anti-CD20) Serotherapy of Human B-Cell Lymphomas." *Blood* 69/2:584-591 (1987). A potential problem with this approach is that non-human monoclonal antibodies (*eg*, murine monoclonal antibodies) typically lack human effector functionality, *ie* they are unable to, *inter alia*, mediate complement dependent lysis or lyse human target cells through antibody dependent cellular toxicity or Fc-receptor mediated phagocytosis. Furthermore, non-human monoclonal antibodies can be recognized by the human host as a foreign protein; therefore, repeated injections of such foreign antibodies can lead to the induction of immune responses leading to harmful hypersensitivity reactions. For murine-based monoclonal antibodies, this is often referred to as a Human Anti-Mouse Antibody response, or "HAMA" response. Additionally, these "foreign" antibodies can be attacked by the immune system of the host such that they are, in effect, neutralized before they reach their target site.

Lymphocytes and lymphoma cells are inherently sensitive to radiotherapy for several reasons: the local emission of ionizing radiation of radiolabeled antibodies may kill cells with or without the target antigen (*eg*, CD20) in close proximity to antibody bound to the antigen; penetrating radiation may obviate the problem of limited access to the antibody in bulky or poorly vascularized tumors; and, the total amount of antibody required may be reduced. The radionuclide emits radioactive particles which can damage cellular DNA to the point where the cellular repair mechanisms are unable to allow the cell to continue living; therefore, if the target cells are tumors, the radioactive label beneficially kills the tumor cells. Radiolabeled antibodies, by definition, include the use of a radioactive substance which may require the need for precautions for both the patient (*ie* possible bone marrow transplantation) as well as the health care provider (*ie* the need to exercise a high degree of caution when working with the radioactivity).

Therefore, an approach at improving the ability of murine monoclonal antibodies to be effective in the treatment of B-cell disorders has been to conjugate a radioactive label or toxin to the antibody such that the label or toxin is localized at the tumor site. For example, the above-referenced 1F5 antibody has been "labeled" with iodine-131 ( $^{131}\text{I}$ ) and was reportedly evaluated for biodistribution in two patients. See Eary, J.F. *et al.*, "Imaging and Treatment of B-Cell Lymphoma" *J. Nuc. Med.* 31/8:1257-1268 (1990); see also, Press, O.W. *et al.*, "Treatment of Refractory Non-Hodgkin's

Lymphoma with Radiolabeled MB-1 (Anti-CD37) Antibody" *J. Clin. Onc.* 7/8:1027-1038 (1989) (indication that one patient treated with <sup>131</sup>I-labeled IF-5 achieved a "partial response"); Goldenberg, D.M. *et al.*, "Targeting, Dosimetry and Radioimmunotherapy of B-Cell Lymphomas with Iodine-131-Labeled LL2 Monoclonal Antibody" *J. Clin. Onc.* 9/4:548-564 (1991)(three of eight patients receiving multiple injections reported to have developed a HAMA response); Appelbaum, F.R. "Radiolabeled Monoclonal Antibodies in the Treatment of Non-Hodgkin's Lymphoma" *Hem. /Onc. Clinics of N.A.* 5/5:1013-1025 (1991) (review article); Press, O.W. *et al* "Radiolabeled-Antibody Therapy of B-Cell Lymphoma with Autologous Bone Marrow Support." *New England Journal of Medicine* 329/17: 1219-12223 (1993) (iodine-131 labeled anti-CD20 antibody IF5 and B1); and Kaminski, M.G. *et al* "Radioimmunotherapy of B-Cell Lymphoma with [<sup>131</sup>I] Anti-B1 (Anti-CD20) Antibody". *NEJM* 329/7 (1993) (iodine-131 labeled anti-CD20 antibody B1; hereinafter "Kaminski").

Toxins (*ie* chemotherapeutic agents such as doxorubicin or mitomycin C) have also been conjugated to antibodies. See, for example, PCT published application WO 92/07466 (published May 14, 1992).

"Chimeric" antibodies, *ie* antibodies which comprise portions from two or more different species (*eg*, mouse and human) have been developed as an alternative to "conjugated" antibodies. For example, Liu, A.Y. *et al.*, "Production of a Mouse-Human Chimeric Monoclonal Antibody to CD20 with Potent Fc-Dependent Biologic Activity" *J. Immun.* 139/10:3521-3526 (1987), describes a mouse/human chimeric antibody directed against the CD20 antigen. See also, PCT Publication No. WO 88/04936. However, no information is provided as to the ability, efficacy or practicality of using such chimeric antibodies for the treatment of B cell disorders in the reference. It is noted that *in vitro* functional assays (*eg* complement dependent lysis ("CDC"); antibody dependent cellular cytotoxicity ("ADCC"), etc.) cannot inherently predict the *in vivo* capability of a chimeric antibody to destroy or deplete target cells expressing the specific antigen. See, for example, Robinson, R.D. *et al.*, "Chimeric mouse-human anti-carcinoma antibodies that mediate different anti-tumor cell biological activities," *Hum. Antibod. Hybridomas* 2:84-93 (1991)(chimeric mouse-human antibody having undetectable ADCC activity). Therefore, the potential therapeutic efficacy of chimeric antibody can only truly be assessed by *in vivo* experimentation.

What is needed, and what would be a great advance in the art, are therapeutic approaches targeting the CD20 antigen for the treatment of B cell lymphomas in primates, including, but not limited to, humans.

### **C. SUMMARY OF THE INVENTION**

Disclosed herein are therapeutic methods designed for the treatment of B cell disorders, and in particular, B cell lymphomas. These protocols are based upon the administration of immunologically active chimeric anti-CD20 antibodies for the depletion of peripheral blood B cells, including B cells associated with lymphoma; administration of radiolabeled anti-CD20 antibodies for targeting localized and peripheral B cell associated tumors; and administration of chimeric anti-CD20 antibodies and radiolabeled anti-CD20 antibodies in a cooperative therapeutic strategy.

### **D. BRIEF DESCRIPTION OF THE DRAWINGS**

Figure 1 is a diagrammatic representation of a tandem chimeric antibody expression vector useful in the production of immunologically active chimeric anti-CD20 antibodies ("TCAE 8");

Figures 2A through 2E are the nucleic acid sequence of the vector of Figure 1;

Figures 3A through 3F are the nucleic acid sequence of the vector of Figure 1 further comprising murine light and heavy chain variable regions ("anti-CD20 in TCAE 8");

Figure 4 is the nucleic acid and amino acid sequences (including CDR and framework regions) of murine variable region light chain derived from murine anti-CD20 monoclonal antibody 2B8;

Figure 5 is the nucleic acid and amino acid sequences (including CDR and framework regions) of murine variable region heavy chain derived from murine anti-CD20 monoclonal antibody 2B8;

Figure 6 are flow cytometry results evidencing binding of fluorescent-labeled human C1q to chimeric anti-CD20 antibody, including, as controls labeled C1q; labeled C1q and murine anti-CD20 monoclonal antibody 2B8; and labeled C1q and human IgG1,k;

Figure 7 represents the results of complement related lysis comparing chimeric anti-CD20 antibody and murine anti-CD20 monoclonal antibody 2B8;

Figure 8 represents the results of antibody mediated cellular cytotoxicity with *in vivo* human effector cells comparing chimeric anti-CD20 antibody and 2B8;

Figure 9A, 9B and 9C provide the results of non-human primate peripheral blood B lymphocyte depletion after infusion of 0.4 mg/kg (A); 1.6 mg/kg (B); and 6.4 mg/kg (C) of immunologically active chimeric anti-CD20 antibody;

Figure 10 provides the results of, *inter alia*, non-human primate peripheral blood B lymphocyte depletion after infusion of 0.01 mg/kg of immunologically active chimeric anti-CD20 antibody;

Figure 11 provides results of the tumoricidal impact of Y2B8 in a mouse xenographic model utilizing a B cell lymphoblastic tumor;

Figure 12 provides results of the tumoricidal impact of C2B8 in a mouse xenographic model utilizing a B cell lymphoblastic tumor;

Figure 13 provides results of the tumoricidal impact of a combination of Y2B8 and C2B8 in a mouse xenographic model utilizing a B cell lymphoblastic tumor; and

Figures 14A and 14B provide results from a Phase I/II clinical analysis of C2B8 evidencing B-cell population depletion over time for patients evidencing a partial remission of the disease (14A) and a minor remission of the disease (14B).

## **E. DETAILED DESCRIPTION OF PREFERRED EMBODIMENTS**

Generally, antibodies are composed of two light chains and two heavy chain molecules; these chains form a general "Y" shape, with both light and heavy chains forming the arms of the Y and the heavy chains forming the base of the Y. Light and heavy chains are divided into domains of structural and functional homology. The variable domains of both the light ("V<sub>L</sub>") and the heavy ("V<sub>H</sub>") chains determine recognition and specificity. The constant region domains of light ("C<sub>L</sub>") and heavy ("C<sub>H</sub>") chains confer important biological properties, *eg* antibody chain association, secretion, transplacental mobility, Fc receptor binding complement binding, etc. The series of events leading to immunoglobulin gene expression in the antibody producing cells are complex. The variable domain region gene sequences are located in separate germ line gene segments referred to as "V<sub>H</sub>," "D," and "J<sub>H</sub>," or "V<sub>L</sub>" and "J<sub>L</sub>." These gene segments are joined by DNA rearrangements to form the complete V regions expressed in heavy and light chains, respectively. The rearranged, joined V segments (V<sub>L</sub>-J<sub>L</sub> and V<sub>H</sub>-D-J<sub>H</sub>) then encode the complete variable regions or antigen binding domains of light and heavy chains, respectively.

Serotherapy of human B cell lymphomas using an anti-CD20 murine monoclonal antibody (1F5) has been described by Press *et al.*, (69 *Blood* 584, 1987, *supra*); the reported therapeutic responses, unfortunately, were transient. Additionally, 25% of the tested patients reportedly developed a human anti-mouse antibody (HAMA) response to the serotherapy. Press *et al.*, suggest that these antibodies, conjugated to toxins or radioisotopes, might afford a more lasting clinical benefit than the unconjugated antibody.

Owing to the debilitating effects of B cell lymphoma and the very real need to provide viable treatment approaches to this disease, we have embarked upon different approaches having a particular antibody, 2B8, as the common link between the approaches. One such approach advantageously exploits the ability of mammalian systems to readily and efficiently recover peripheral blood B cells; using this approach, we seek to, in essence, purge or deplete B cells in peripheral blood and lymphatic tissue as a means of also removing B cell lymphomas. We accomplish this by utilization of, *inter alia*, immunologically active, chimeric anti-CD20 antibodies. In another approach, we seek to target tumor cells for destruction with radioactive labels.

As used herein, the term "anti-CD20 antibody" is an antibody which specifically recognizes a cell surface non-glycosylated phosphoprotein of 35,000 Daltons, typically designated as the human B lymphocyte restricted differentiation antigen Bp35, commonly referred to as CD20. As used herein, the term "chimeric" when used in reference to anti-CD20 antibodies, encompasses antibodies which are most preferably derived using recombinant deoxyribonucleic acid techniques and which comprise both human (including immunologically "related" species, *eg*, chimpanzee) and non-human components: the constant region of the chimeric antibody is most preferably substantially identical to the constant region of a natural human antibody; the variable region of the chimeric antibody is most preferably derived from a non-human source and has the desired antigenic and specificity to the CD20 cell surface antigen. The non-human source can be any vertebrate source which can be used to generate antibodies to a human CD20 cell surface antigen or material comprising a human CD20 cell surface antigen. Such non-human source includes, but is not limited to, rodents (*eg*, rabbit, rat, mouse, etc.) and non-human primates (*eg*, Old World Monkey, Ape, etc.). Most preferably, the non-human component (variable region) is derived from a murine source. As used herein, the phrase "immunologically active" when used in reference to chimeric anti-CD20 antibodies, means a chimeric antibody which binds human C1q, mediates complement dependent lysis ("CDC") of human B lymphoid cell lines, and lyses human target cells through antibody dependent cellular cytotoxicity ("ADCC"). As used herein, the phrases "indirect labeling" and "indirect labeling

approach" both mean that a chelating agent is covalently attached to an antibody and at least one radionuclide is inserted into the chelating agent. Preferred chelating agents and radionuclides are set forth in Srivagtava, S.C. and Mease, R.C., "Progress in Research on Ligands, Nuclides and Techniques for Labeling Monoclonal Antibodies," *Nucl. Med. Bio.* 18/6: 589-603 (1991) ("Srivagtava") which is incorporated herein by reference. A particularly preferred chelating agent is 1-isothiocyanatobenzyl-3-methyldiothelene triaminepent acetic acid ("MX-DTPA"); particularly preferred radionuclides for indirect labeling include indium [111] and yttrium [90]. As used herein, the phrases "direct labeling" and "direct labeling approach" both mean that a radionuclide is covalently attached directly to an antibody (typically via an amino acid residue). Preferred radionuclides are provided in Srivagtava; a particularly preferred radionuclide for direct labeling is iodine [131] covalently attached via tyrosine residues. The indirect labeling approach is particularly preferred.

The therapeutic approaches disclosed herein are based upon the ability of the immune system of primates to rapidly recover, or rejuvenate, peripheral blood B cells. Additionally, because the principal immune response of primates is occasioned by T cells, when the immune system has a peripheral blood B cell deficiency, the need for "extraordinary" precautions (*ie* patient isolation, etc.) is not necessary. As a result of these and other nuances of the immune systems of primates, our therapeutic approach to B cell disorders allows for the purging of peripheral blood B cells using immunologically active chimeric anti-CD20 antibodies.

Because peripheral blood B cell disorders, by definition, can indicate a necessity for access to the blood for treatment, the route of administration of the immunologically active chimeric anti-CD20 antibodies and radio-labeled anti-CD20 antibodies is preferably parenteral; as used herein, the term "parenteral" includes intravenous, intramuscular, subcutaneous, rectal, vaginal or intraperitoneal administration. Of these, intravenous administration is most preferred.

The immunologically active chimeric anti-CD20 antibodies and radiolabeled anti-CD20 antibodies will typically be provided by standard technique within a pharmaceutically acceptable buffer, for example, sterile saline, sterile buffered water, propylene glycol, combinations of the foregoing, etc. Methods for preparing parenterally administerable agents are described in *Pharmaceutical Carriers & Formulations*, Martin, Remington's Pharmaceutical Sciences, 15th Ed. (Mack Pub. Co., Easton, PA 1975), which is incorporated herein by reference.

The specific, therapeutically effective amount of immunologically active chimeric anti-CD20 antibodies useful to produce a unique therapeutic effect in any given patient can be determined by standard techniques well known to those of ordinary skill in the art.

Effective dosages (*ie* therapeutically effective amounts) of the immunologically active chimeric anti-CD20 antibodies range from about 0.001 to about 30 mg/kg body weight, more preferably from about 0.01 to about 25 mg/kg body weight, and most preferably from about 0.4 to about 20.0 mg/kg body weight. Other dosages are viable; factors influencing dosage include, but are not limited to, the severity of the disease; previous treatment approaches; overall health of the patient; other diseases present, etc. The skilled artisan is readily credited with assessing a particular patient and determining a suitable dosage that falls within the ranges, or if necessary, outside of the ranges.

Introduction of the immunologically active chimeric anti-CD20 antibodies in these dose ranges can be carried out as a single treatment or over a series of treatments. With respect to chimeric antibodies, it is preferred that such introduction be carried out over a series of treatments; this preferred approach is predicated upon the treatment methodology associated with this disease. While not wishing to be bound by any particular theory, because the immunologically active chimeric anti-CD20 antibodies are both immunologically active and bind to CD20, upon initial introduction of the immunologically active chimeric anti-CD20 antibodies to the individual, peripheral blood B cell depletion will begin; we have observed a nearly complete depletion within about 24 hours post treatment infusion. Because of this, subsequent introduction(s) of the immunologically active chimeric anti-CD20 antibodies (or radiolabeled anti-CD20 antibodies) to the patient is presumed to: a) clear remaining peripheral blood B cells; b) begin B cell depletion from lymph nodes; c) begin B cell depletion from other tissue sources, *eg*, bone marrow, tumor, etc. Stated again, by using repeated introductions of the immunologically active chimeric anti-CD20 antibodies, a series of events take place, each event being viewed by us as important to effective treatment of the disease. The first "event" then, can be viewed as principally directed to substantially depleting the patient's peripheral blood B cells; the subsequent "events" can be viewed as either principally directed to simultaneously or serially clearing remaining B cells from the system clearing lymph node B cells, or clearing other tissue B cells.

In effect, while a single dosage provides benefits and can be effectively utilized for disease treatment/management, a preferred treatment course can occur over several stages; most preferably, between about 0.4 and about 20 mg/kg body weight of the immunologically active chimeric anti-CD20 antibodies is introduced to the patient once a week for between about 2 to 10 weeks, most preferably for about 4 weeks.

With reference to the use of radiolabeled anti-CD20 antibodies, a preference is that the antibody is non-chimeric; this preference is predicted upon the significantly longer circulating half-life of chimeric antibodies vis-a-vis murine antibodies (*ie* with a longer circulating half-life, the radionuclide is present in the patient for extended periods). However, radiolabeled chimeric antibodies can be beneficially utilized with lower milli-Curies ("mCi") dosages used in conjunction with the chimeric antibody relative to the murine antibody. This scenario allows for a decrease in bone marrow toxicity to an acceptable level, while maintaining therapeutic utility.



A variety of radionuclides are applicable to the present invention and those skilled in the art are credited with the ability to readily determine which radionuclide is most appropriate under a variety of circumstances. For example, iodine [131] is a well known radionuclide used for targeted immunotherapy. However, the clinical usefulness of iodine [131] can be limited by several factors including: eight-day physical half-life; dehalogenation of iodinated antibody both in the blood and at tumor sites; and emission characteristics (*eg* large gamma component) which can be suboptimal for localized dose deposition in tumor. With the advent of superior chelating agents, the opportunity for attaching metal chelating groups to proteins has increased the opportunities to utilize other radionuclides such as indium [131] and yttrium [90]. Yttrium [90] provides several benefits for utilization in radioimmunotherapeutic applications: the 64 hour half-life of yttrium [90] is long enough to allow antibody accumulation by tumor and, unlike *eg* iodine [131], yttrium [90] is a pure beta emitter of high energy with no accompanying gamma irradiation in its decay, with a range in tissue of 100 to 1000 cell diameters. Furthermore, the minimal amount of penetrating radiation allows for outpatient administration of yttrium [90]-labeled antibodies. Furthermore, internalization of labeled antibody is not required for cell killing, and the local emission of ionizing radiation should be lethal for adjacent tumor cells lacking the target antigen.

One non-therapeutic limitation to yttrium [90] is based upon the absence of significant gamma radiation making imaging therewith difficult. To avoid this problem, a diagnostic "imaging" radionuclide, such as indium [111], can be utilized for determining the location and relative size of a tumor prior to the administration of therapeutic doses of yttrium [90]-labeled anti-CD20. Indium [111] is particularly preferred as the diagnostic radionuclide because: between about 1 to about 10mCi can be safely administered without detectable toxicity; and the imaging data is generally predictive of subsequent yttrium [90]-labeled antibody distribution. Most imaging studies utilize 5mCi indium [111]-labeled antibody because this dose is both safe and has increased imaging efficiency compared with lower doses, with optimal imaging occurring at three to six days after antibody administration. *See, for example*, Murray J.L., 26 *J. Nuc. Med.* 3328 (1985) and Carraguallo, J.A. et al., 26 *J. Nuc. Med.* 67 (1985).

Effective single treatment dosages (*ie* therapeutically effective amounts) of yttrium [90] labeled anti-CD20 antibodies range from between about 5 and about 75mCi, more preferably between about 10 and about 40mCi. Effective single treatment non-marrow ablative dosages of iodine [131] labeled anti-CD20 antibodies range from between about 5 and about 70mCi, more preferably between about 5 and about 40mCi. Effective single treatment ablative dosages (*ie* may require autologous bone marrow transplantation) of iodine [131] labeled anti-CD20 antibodies range from between about 30 and about 600mCi, more preferably between about 50 and less than about 500mCi. In conjunction with a chimeric anti-CD20 antibody, owing to the longer circulating half life vis-a-vis murine antibodies, an effective single treatment non-marrow ablative dosages of iodine [131] labeled chimeric anti-CD20 antibodies range from between about 5 and about 40mCi, more preferably less than about 30mCi. Imaging criteria for, *eg* the indium [111] label, are typically less than about 5mCi.

With respect to radiolabeled anti-CD20 antibodies, therapy therewith can also occur using a single therapy treatment or using multiple treatments. Because of the radionuclide component, it is preferred that prior to treatment, peripheral stem cells ("PSC") or bone marrow ("BM") be "harvested" for patients experiencing potentially fatal bone marrow toxicity resulting from radiation. BM and/or PSC are harvested using standard techniques, and then purged and frozen for possible reinfusion. Additionally, it is most preferred that prior to treatment a diagnostic dosimetry study using a diagnostic labeled antibody (*eg* using indium [111]) be conducted on the patient, a purpose of which is to ensure that the therapeutically labeled antibody (*eg* using yttrium [90]) will not become unnecessarily "concentrated" in any normal organ or tissue.

Chimeric mouse/human antibodies have been described. *See, for example*, Morrison, S.L. et al., *PNAS* 11:6851-6854 (November 1984); European Patent Publication No. 173494; Boulianne, G.L. et al., *Nature* 312:643 (December 1984); Neubeiger, M.S. et al., *Nature* 314:268 (March 1985); European Patent Publication No. 125023; Tan et al., *J. Immunol.* 135:8564 (November 1985); Sun, L.K. et al., *Hybridoma* 5/1:517 (1986); Sahagan et al., *J. Immunol.* 137:1066-1074 (1986). *See generally*, Muron, *Nature* 312:597 (December 1984); Dickson, *Genetic Engineering News* 5/3 (March 1985); Marx, *Science* 229 455 (August 1985); and Morrison *Science* 229:1202-1207 (September 1985). Robinson et al., in PCT Publication Number WO 88/04936 describe a chimeric antibody with human constant region and murine variable region, having specificity to an epitope of CD20; the murine portion of the chimeric antibody of the Robinson references is derived from the 2H7 mouse monoclonal antibody (gamma 2b, kappa). While the reference notes that the described chimeric antibody is a "prime candidate" for the treatment of B cell disorders, this statement can be viewed as no more than a suggestion to those in the art to determine whether or not this suggestion is accurate for this particular antibody, particularly because the reference lacks any data to support an assertion of therapeutic effectiveness, and importantly, data using higher order mammals such as primates or humans.

Methodologies for generating chimeric antibodies are available to those in the art. For example, the light and heavy chains can be expressed separately, using, for example, immunoglobulin light chain and immunoglobulin heavy chains in separate plasmids. These can then be purified and assembled *in vitro* into complete antibodies; methodologies for accomplishing such assembly have been described. *See, for example*, Scharff, M., *Harvey Lectures* 69:125 (1974). *In vitro* reaction parameters for the formation of IgG antibodies from reduced isolated light and heavy chains have also been described. *See, for example*, Beychok, S., *Cells of Immunoglobulin Synthesis*, Academic Press, New York, p. 69,

1979. Co-expression of light and heavy chains in the same cells to achieve intracellular association and linkage of heavy and light chains into complete H<sub>2</sub>L<sub>2</sub> IgG antibodies is also possible. Such co-expression can be accomplished using either the same or different plasmids in the same host cell.

Another approach, and one which is our most preferred approach for developing a chimeric non-human/human anti-CD20 antibody, is based upon utilization of an expression vector which includes, *ab initio*, DNA encoding heavy and light chain constant regions from a human source. Such a vector allows for inserting DNA encoding non-human variable region such that a variety of non-human anti-CD20 antibodies can be generated, screened and analyzed for various characteristics (*eg* type of binding specificity, epitope binding regions, etc.); thereafter, cDNA encoding the light and heavy chain variable regions from a preferred or desired anti-CD20 antibody can be incorporated into the vector. We refer to these types of vectors as Tandem Chimeric Antibody Expression ("TCAE") vectors. A most preferred TCAE vector which was used to generate immunologically active chimeric anti-CD20 antibodies for therapeutic treatment of lymphomas is TCAE 8. TCAE 8 is a derivative of a vector owned by the assignee of this patent document, referred to as TCAE 5.2 the difference being that in TCAE 5.2, the translation initiation start site of the dominant selectable marker (neomycin phosphotransferase, "NEO") is a consensus Kozak sequence, while for TCAE 8, this region is a partially impaired consensus Kozak sequence. Details regarding the impact of the initiation start site of the dominant selectable marker of the TCAE vectors (also referred to as "ANEX vector") vis-a-vis protein expression are disclosed in detail in the co-pending application filed herewith.

TCAE 8 comprises four (4) transcriptional cassettes, and these are in tandem order, *ie* a human immunoglobulin light chain absent a variable region; a human immunoglobulin heavy chain absent a variable region; DHFR; and NEO. Each transcriptional cassette contains its own eukaryotic promotor and polyadenylation region (reference is made to Figure 1 which is a diagrammatic representation of the TCAE 8 vector). Specifically:

1) the CMV promoter/enhancer in front of the immunoglobulin heavy chain is a truncated version of the promoter/enhancer in front of the light chain, from the Nhe I site at -350 to the Sst I site at -16 (*see*, 41 *Cell* 521, 1985).

2) a human immunoglobulin light chain constant region was derived via amplification of cDNA by a PCR reaction. In TCAE 8, this was the human immunoglobulin light chain kappa constant region (Kabat numbering, amino acids 108-214, allotype Km 3, (*see*, Kabat, E.A. "Sequences of proteins of immunological interest," NIH Publication, Fifth Ed. No. 91-3242, 1991)), and the human immunoglobulin heavy chain gamma 1 constant region (Kabat numbering amino acids 114-478, allotype Gmla, Gmlz). The light chain was isolated from normal human blood (IDEC Pharmaceuticals Corporation, La Jolla, CA); RNA therefrom was used to synthesize cDNA which was then amplified using PCR techniques (primers were derived vis-a-vis the consensus from Kabat). The heavy chain was isolated (using PCR techniques) from cDNA prepared from RNA which was in turn derived from cells transfected with a human IgG1 vector (*see*, 3 *Prot. Eng.* 531, 1990; vector pN<sub>1</sub>62). Two amino acids were changed in the isolated human IgG1 to match the consensus amino acid sequence from Kabat, to wit: amino acid 225 was changed from valine to alanine (GTT to GCA), and amino acid 287 was changed from methionine to lysine (ATG to AAG);

3) The human immunoglobulin light and heavy chain cassettes contain synthetic signal sequences for secretion of the immunoglobulin chains;

4) The human immunoglobulin light and heavy chain cassettes contain specific DNA restriction sites which allow for insertion of light and heavy immunoglobulin variable regions which maintain the transitional reading frame and do not alter the amino acids normally found in immunoglobulin chains;

5) The DHFR cassette contained its own eukaryotic promoter (mouse beta globin major promoter, "BETA") and polyadenylation region (bovine growth hormone polyadenylation, "BGH"); and

6) The NEO cassette contained its own eukaryotic promoter (BETA) and polyadenylation region (SV40 early polyadenylation, "SV").

With respect to the TCAE 8 vector and the NEO cassette, the Kozak region was a partially impaired consensus Kozak sequence (which included an upstream Cla I site):

ClaI      -3      +1  
 GGGAGCTTGG ATCGAT ccTct ATG Gtt

(In the TCAE 5.2 vector, the change is between the ClaI and ATG regions, to wit: ccAcc.)

The complete sequence listing of TCAE 8 (including the specific components of these four transcriptional cassettes) is set forth in Figure 2 (SEQ. ID. NO. 1).

As will be appreciated by those in the art, the TCAE vectors beneficially allow for substantially reducing the time in generating the immunologically active chimeric anti-CD20 antibodies. Generation and isolation of non-human light and heavy chain variable regions, followed by incorporation thereof within the human light chain constant transcriptional cassette and human heavy chain constant transcriptional cassette, allows for production of immunologically active chimeric anti-CD20 antibodies.

We have derived a most preferred non-human variable region with specificity to the CD20 antigen using a murine source and hybridoma technology. Using polymerase chain reaction ("PCR") techniques, the murine light and heavy variable regions were cloned directly into the TCAE 8 vector--this is the most preferred route for incorporation of the non-human variable region into the TCAE vector. This preference is principally predicated upon the efficiency of the PCR reaction and the accuracy of insertion. However, other equivalent procedures for accomplishing this task are available. For example, using TCAE 8 (or an equivalent vector), the sequence of the variable region of a non-human anti-CD20 antibody can be obtained, followed by oligonucleotide synthesis of portions of the sequence or, if appropriate, the entire sequence; thereafter, the portions or the entire synthetic sequence can be inserted into the appropriate locations within the vector. Those skilled in the art are credited with the ability to accomplish this task.

Our most preferred immunologically active chimeric anti-CD20 antibodies were derived from utilization of TCAE 8 vector which included murine variable regions derived from monoclonal antibody to CD20; this antibody (to be discussed in detail, *infra*), is referred to as "2B8." The complete sequence of the variable regions obtained from 2B8, in TCAE 8 ("anti-CD20 in TCAE 8") is set forth in Figure 3 (SEQ. ID. NO. 2).

The host cell line utilized for protein expression is most preferably of mammalian origin; those skilled in the art are credited with ability to preferentially determine particular host cell lines which are best suited for the desired gene product to be expressed therein. Exemplary host cell lines include, but are not limited to, DG44 and DUXBII (Chinese Hamster Ovary lines, DHFR minus), HELA (human cervical carcinoma), CVI (monkey kidney line), COS (a derivative of CVI with SV40 T antigen), R1610 (Chinese hamster fibroblast) BALBC/3T3 (mouse fibroblast), HAK (hamster kidney line), SP2/O (mouse myeloma), P3x63-Ag3.653 (mouse myeloma), BFA-IcIBPT (bovine endothelial cells), RAJI (human lymphocyte) and 293 (human kidney). Host cell lines are typically available from commercial services, the American Tissue Culture Collection or from published literature.

Preferably the host cell line is either DG44 ("CHO") or SP2/O. See Urland, G. *et al.*, "Effect of gamma rays and the dihydrofolate reductase locus: deletions and inversions." *Som. Cell & Mol. Gen.* 12/6:555-566 (1986), and Shulman, M. *et al.*, "A better cell line for making hybridomas secreting specific antibodies." *Nature* 276:269 (1978), respectively. Most preferably, the host cell line is DG44. Transfection of the plasmid into the host cell can be accomplished by any technique available to those in the art. These include, but are not limited to, transfection (including electrophoresis and electroporation), cell fusion with enveloped DNA, microinjection, and infection with intact virus. See, Ridgway, A.A.G. "Mammalian Expression Vectors." Chapter 24.2, pp. 470-472 *Vectors*, Rodriguez and Denhardt, Eds. (Butterworths, Boston, MA 1988). Most preferably, plasmid introduction into the host is via electroporation.

## F. EXAMPLES

The following examples are not intended, nor are they to be construed as limiting the invention. The examples are intended to evidence: dose-imaging using a radiolabeled anti-CD20 antibody ("I2B8"); radiolabeled anti-CD20 antibody ("Y2B8"); and immunologically active, chimeric anti-CD20 antibody ("C2B8") derived utilizing a specific vector ("TCAE 8") and variable regions derived from murine anti-CD20 monoclonal antibody ("2B8").

### I. RADIOLABELED ANTI-CD20 ANTIBODY 2B8

#### A. *Anti-CD20 Monoclonal Antibody (Murine) Production ("2B8")*

BALB/C mice were repeatedly immunized with the human lymphoblastoid cell line SB (see, Adams, R.A. *et al.*, "Direct implantation and serial transplantation of human acute lymphoblastic leukemia in hamsters, SB-2." *Can Res* 28:1121-1125 (1968); this cell line is available from the American Tissue Culture Collection, Rockville, MD., under ATCC accession number ATCC CCL 120), with weekly injections over a period of 3-4 months. Mice evidencing high serum titers of anti-CD20 antibodies, as determined by inhibition of known CD20-specific antibodies (anti-CD20 antibodies utilized were Leu 16, Beckton Dickinson, San Jose, CA, Cat. No. 7670; and BI, Coulter Corp., Hialeah, FL, Cat. No. 6602201) were identified; the spleens of such mice were then removed. Spleen cells were fused with the mouse myeloma SP2/O in accordance with the protocol described in Einfeld, D.A. *et al.*, (1988) *EMBO* 7:711 (SP2/O has ATCC accession no. ATCC CRL 8006).

Assays for CD20 specificity were accomplished by radioimmunoassay. Briefly, purified anti-CD20 BI was radiola-

beled with  $I^{125}$  by the iodobead method as described in Valentine, M.A. *et al.*, (1989) *J. Biol. Chem.* 264:11282. ( $I^{125}$  Sodium Iodide, ICN, Irvine, CA, Cat. No. 28665H). Hybridomas were screened by co-incubation of 0.05 ml of media from each of the fusion wells together with 0.05 ml of  $I^{125}$  labeled anti-CD20 BI (10 ng) in 1% BSA, PBS (pH 7.4), and 0.5 ml of the same buffer containing 100,000 SB cells. After incubation for 1 hr at room temperature, the cells were harvested by transferring to 96 well titer plates (V&P Scientific, San Diego, CA), and washed thoroughly. Duplicate wells containing unlabeled anti-CD20 BI and wells containing no inhibiting antibody were used as positive and negative controls, respectively. Wells containing greater than 50% inhibition were expanded and cloned. The antibody demonstrating the highest inhibition was derived from the cloned cell line designated herein as "2B8."

#### B. Preparation of 2B8-MX-DTPA Conjugate

##### i. MX-DTPA

Carbon-14-labeled 1-isothiocyanatobenzyl-3-methyldiethylene triaminepentaacetic acid ("carbon-14 labeled MX-DTPA") was used as a chelating agent for conjugation of radiolabel to 2B8. Manipulations of MX-DTPA were conducted to maintain metal-free conditions, *ie* metal-free reagents were utilized and, when possible, polypropylene plastic containers (flasks, beakers, graduated cylinders, pipette tips) washed with Alconox and rinsed with Milli-Q water, were similarly utilized. MX-DTPA was obtained as a dry solid from Dr. Otto Gansow (National Institute of Health, Bethesda, MD) and stored desiccated at 4°C (protected from light), with stock solutions being prepared in Milli-Q water at a concentration of 2-5mM, with storage at -70°C. MX-DTPA was also obtained from Coulter Immunology (Hialeah, Florida) as the disodium salt in water and stored at -70°C.

##### ii. Preparation of 2B8

Purified 2B8 was prepared for conjugation with MX-DTPA by transferring the antibody into metal-free 50mM bicine-NaOff, pH 8.6, containing 150 mM NaCl, using repetitive buffer exchange with CENTRICON 30™ spin filters (30,000D, MWCO; Amicon). Generally, 50-200  $\mu$ L of protein (10 mg/ml) was added to the filter unit, followed by 2 mL of bicine buffer. The filter was centrifuged at 4°C in a Sorval SS-34 rotor (6,000 rpm, 45 min.). Retentate volume was approximately 50-100  $\mu$ L; this process was repeated twice using the same filter. Retentate was transferred to a polypropylene 1.5 mL screw cap tube, assayed for protein, diluted to 10.0 mg/mL and stored at 4°C until utilized; protein was similarly transferred into 50 mM sodium citrate, pH 5.5, containing 150 mM NaCl and 0.05% sodium azide, using the foregoing protocol.

##### iii. Coniugation of 2B8 with MX-DTPA

Conjugation of 2B8 with MX-DTPA was performed in polypropylene tubes at ambient temperature. Frozen MX-DTPA stock solutions were thawed immediately prior to use. 50-200  $\mu$ L of protein at 10 mg/mL were reacted with MX-DTPA at a molar ratio of MX-DTPA-to-2B8 of 4:1. Reactions were initiated by adding the MX-DTPA stock solution and gently mixing; the conjugation was allowed to proceed overnight (14 to 20 hr), at ambient temperature. Unreacted MX-DTPA was removed from the conjugate by dialysis or repetitive ultrafiltration, as described above in Example I.B.ii, into metal-free normal saline (0.9% w/v) containing 0.05% sodium azide. The protein concentration was adjusted to 10 mg/mL and stored at 4°C in a polypropylene tube until radiolabeled.

##### iv. Determination of MX-DTPA Incorporation

MX-DTPA incorporation was determined by scintillation counting and comparing the value obtained with the purified conjugate to the specific activity of the carbon-[14]-labeled MX-DTPA. For certain studies, in which non-radioactive MX-DTPA (Coulter Immunology) was utilized, MX-DTPA incorporation was assessed by incubating the conjugate with an excess of a radioactive carrier solution of yttrium-[90] of known concentration and specific activity.

A stock solution of yttrium chloride of known concentration was prepared in metal-free 0.05 N HCl to which carrier-free yttrium-[90] (chloride salt) was added. An aliquot of this solution was analyzed by liquid scintillation counting to determine an accurate specific activity for this reagent. A volume of the yttrium chloride reagent equal to 3-times the number of mols of chelate expected to be attached to the antibody, (typically 2 mol/mol antibody), was added to a polypropylene tube, and the pH adjusted to 4.0-4.5 with 2 M sodium acetate. Conjugated antibody was subsequently added and the mixture incubated 15-30 min. at ambient temperature. The reaction was quenched by adding 20 mM EDTA to a final concentration of 1 mM and the pH of the solution adjusted to approximately pH 6 with 2M sodium acetate.

After a 5 min. incubation, the entire volume was purified by high-performance, size-exclusion chromatography (described *infra*). The eluted protein-containing fractions were combined, the protein concentration determined, and an aliquot assayed for radioactivity. The chelate incorporation was calculated using the specific activity of the yttrium-[90]

chloride preparation and the protein concentration.

#### v. Immunoreactivity of 2B8-MX-DTPA

The immunoreactivity of conjugated 2B8 was assessed using whole-cell ELISA. Mid-log phase SB cells were harvested from culture by centrifugation and washed two times with 1X HBSS. Cells were diluted to  $1-2 \times 10^6$  cells/mL in HBSS and aliquoted into 96-well polystyrene microtiter plates at 50,000- 100,000 cells/well. The plates were dried under vacuum for 2 h. at 40-45°C to fix the cells to the plastic; plates were stored dry at -20°C until utilized. For assay, the plates were warmed to ambient temperature immediately before use, then blocked with 1X PBS, pH 7.2-7.4 containing 1% BSA (2 h). Samples for assay were diluted in 1X PBS/1% BSA, applied to plates and serially diluted (1:2) into the same buffer. After incubating plates for 1 h. at ambient temperature, the plates were washed three times with 1X PBS. Secondary antibody (goat anti-mouse IgG1-specific HRP conjugate 50  $\mu$ L) was added to wells (1:1500 dilution in 1X PBS/1% BSA) and incubated 1 h. at ambient temperature. Plates were washed four times with 1X PBS followed by the addition of ABTS substrate solution (50 mM sodium citrate, pH 4.5 containing 0.01% ATBS and 0.001%  $H_2O_2$ ). Plates were read at 405 nm after 15-30 min. incubation. Antigen-negative HSB cells were included in assays to monitor non-specific binding. Immunoreactivity of the conjugate was calculated by plotting the absorbance values vs. the respective dilution factor and comparing these to values obtained using native antibody (representing 100% immunoreactivity) tested on the same plate; several values on the linear portion of the titration profile were compared and a mean value determined (data not shown).

#### vi. Preparation of Indium-[111]-Labeled 2B8-MX-DTPA ("I2B8")

Conjugates were radiolabeled with carrier-free indium-[111]. An aliquot of isotope (0.1-2 mCi/mg antibody) in 0.05 M HCL was transferred to a polypropylene tube and approximately one-tenth volume of metal-free 2 M HCl added. After incubation for 5 min., metal-free 2 M sodium acetate was added to adjust the solution to pH 4.0-4.4. Approximately 0.5 mg of 2B8-MX-DTPA was added from a stock solution of 10.0 mg/mL DTPA in normal saline, or 50 mM sodium citrate/150 mM NaCl containing 0.05% sodium azide, and the solution gently mixed immediately. The pH solution was checked with pH paper to verify a value of 4.0-4.5 and the mixture incubated at ambient temperature for 15-30 min. Subsequently, the reaction was quenched by adding 20 mM EDTA to a final concentration of 1 mM and the reaction mixture was adjusted to approximately pH 6.0 using 2 M sodium acetate.

After a 5-10 min. incubation, uncomplexed radioisotope was removed by size-exclusion chromatography. The HPLC unit consisted of Waters Model 6000 or TosoHaas Model TSK-6110 solvent delivery system fitted, respectively, with a Waters U6K or Rheodyne 700 injection valve. Chromatographic separations were performed using a gel permeation column (BioRad SEC-250; 7.5 x 300 mm or comparable TosoHaas column) and a SEC-250 guard column (7.5 x 100 mm). The system was equipped with a fraction collector (Pharmacia Frac200) and a UV monitor fitted with a 280 nm filter (Pharmacia model UV-1). Samples were applied and eluted isocratically using 1X PBS, pH 7.4, at 1.0 mL/min flow rate. One-half milliliter fractions were collected in glass tubes and aliquots of these counted in a gamma counter. The lower and upper windows were set to 100 and 500 KeV respectively.

The radioincorporation was calculated by summing the radioactivity associated with the eluted protein peak and dividing this number by the total radioactivity eluted from the column; this value was then expressed as a percentage (data not shown). In some cases, the radioincorporation was determined using instant thin-layer chromatography ("ITLC"). Radiolabeled conjugate was diluted 1:10 or 1:20 in 1X PBS containing or 1X PBS/1 mM DTPA, then 1  $\mu$ L was spotted 1.5 cm from one end of a 1 x 5 cm strip of ITLC SG paper. The paper was developed by ascending chromatography using 10% ammonium acetate in methanol:water (1:1;v/v). The strip was dried, cut in half crosswise, and the radioactivity associated with each section determined by gamma counting. The radioactivity associated with the bottom half of the strip (protein-associated radioactivity) was expressed as a percentage of the total radioactivity, determined by summing the values for both top and bottom halves (data not shown).

Specific activities were determined by measuring the radioactivity of an appropriate aliquot of the radiolabeled conjugate. This value was corrected for the counter efficiency (typically 75%) and related to the protein concentration of the conjugate, previously determined by absorbance at 280 nm, and the resulting value expressed as mCi/mg protein.

For some experiments, 2B8-MX-DTPA was radiolabeled with indium [111] following a protocol similar to the one described above but without purification by HPLC; this was referred to as the "mix-and-shoot" protocol.

#### vii. Preparation of Yttrium-[90]-Labeled 2B8-MX-DTPA ("Y2B8")

The same protocol described for the preparation of I2B8 was followed for the preparation of the yttrium-[90]-labeled 2B8-MX-DTPA ("Y2B8") conjugate except that 2 ng HCl was not utilized; all preparations of yttrium-labeled conjugates were purified by size-exclusion chromatography as described above.

### C. Non-Human Animal Studies.

#### i. Biodistribution of Radiolabeled 2B8-MX-DTPA

I2B8 was evaluated for tissue biodistribution in six-to-eight week old BALB/c mice. The radiolabeled conjugate was prepared using clinical-grade 2B8-MX-DTPA following the "mix and shoot" protocol described above. The specific activity of the conjugate was 2.3 mCi/mg and the conjugate was formulated in PBS, pH 7.4 containing 50mg/mL HSA. Mice were injected intravenously with 100  $\mu$ L of I2B8 (approximately 21  $\mu$ Ci) and groups of three mice were sacrificed by cervical dislocation at 0, 24, 48, and 72 hours. After sacrifice, the tail, heart, lungs, liver, kidney, spleen, muscle, and femur were removed, washed and weighed; a sample of blood was also removed for analysis. Radioactivity associated with each specimen was determined by gamma counting and the percent injected dose per gram tissue subsequently determined. No attempt was made to discount the activity contribution represented by the blood associated with individual organs.

In a separate protocol, aliquots of 2B8-MX-DTPA incubated at 4°C and 30°C for 10 weeks were radiolabeled with indium-[111] to a specific activity of 2.1 mCi/mg for both preparations. These conjugates were then used in biodistribution studies in mice as described above.

For dosimetry determinations, 2B8-MX-DTPA was radiolabeled with indium-[111] to a specific activity of 2.3 mCi/mg and approximately 1.1  $\mu$ Ci was injected into each of 20 BALB/c mice. Subsequently, groups of five mice each were sacrificed at 1, 24, 48 and 72 hours and their organs removed and prepared for analysis. In addition, portions of the skin, muscle and bone were removed and processed for analysis; the urine and feces were also collected and analyzed for the 24-72 hour time points.

Using a similar approach, 2B8-MX-DTPA was also radiolabeled with yttrium-[90] and its biological distribution evaluated in BALB/c mice over a 72-hour time period. Following purification by HPLC size exclusion chromatography, four groups of five mice each were injected intravenously with approximately 1  $\mu$ Ci of clinically-formulated conjugate (specific activity: 12.2 mCi/mg); groups were subsequently sacrificed at 1, 24, 48 and 72 hours and their organs and tissues analyzed as described above. Radioactivity associated with each tissue specimen was determined by measuring bremsstrahlung energy with a gamma scintillation counter. Activity values were subsequently expressed as percent injected dose per gram tissue or percent injected dose per organ. While organs and other tissues were rinsed repeatedly to remove superficial blood, the organs were not perfused. Thus, organ activity values were not discounted for the activity contribution represented by internally associated blood.

#### ii. Tumor Localization of I2B8

The localization of radiolabeled 2B8-MX-DTPA was determined in athymic mice bearing Ramos B cell tumors. Six-to-eight week old athymic mice were injected subcutaneously (left-rear flank) with 0.1 mL of RPMI-1640 containing  $1.2 \times 10^7$  Ramos tumor cells which had been previously adapted for growth in athymic mice. Tumors arose within two weeks and ranged in weight from 0.07 to 1.1 grams. Mice were injected intravenously with 100  $\mu$ L of indium-[111]-labeled 2B8-MX-DTPA (16.7  $\mu$ Ci) and groups of three mice were sacrificed by cervical dislocation at 0, 24, 48, and 72 hours. After sacrifice the tail, heart, lungs, liver, kidney, spleen, muscle, femur, and tumor were removed, washed, weighed; a sample of blood was also removed for analysis. Radioactivity associated with each specimen was determined by gamma counting and the percent injected dose per gram tissue determined.

#### iii. Biodistribution and Tumor Localization Studies with Radiolabeled 2B8-MX-DTPA.

Following the preliminary biodistribution experiment described above (Example I.B.viii.a.), conjugated 2B8 was radiolabeled with indium-[111] to a specific activity of 2.3 mCi/mg and roughly 1.1  $\mu$ Ci was injected into each of twenty BALB/c mice to determine biodistribution of the radiolabeled material. Subsequently, groups of five mice each were sacrificed at 1, 24, 48 and 72 hours and their organs and a portion of the skin, muscle and bone were removed and processed for analysis. In addition, the urine and feces were collected and analyzed for the 24-72 hour time-points. The level of radioactivity in the blood dropped from 40.3% of the injected dose per gram at 1 hour to 18.9% at 72 hours (data not shown). Values for the heart, kidney, muscle and spleen remained in the range of 0.7-9.8% throughout the experiment. Levels of radioactivity found in the lung decreased from 14.2% at 1 hour to 7.6% at 72 hours; similarly the respective liver injected-dose per gram values were 10.3% and 9.9%. These data were used in determining radiation absorbed dose estimates I2B8 described below.

The biodistribution of yttrium-[90]-labeled conjugate, having a specific activity of 12.2 mCi/mg antibody, was evaluated in BALB/c mice. Radioincorporations of >90% were obtained and the radiolabeled antibody was purified by HPLC. Tissue deposition of radioactivity was evaluated in the major organs, and the skin, muscle, bone, and urine and feces over 72 hours and expressed as percent injected dose/g tissue. Results (not shown) evidenced that while the levels of radioactivity associated with the blood dropped from approximately 39.2% injected dose per gram at 1 hour to roughly

15.4% after 72 hours the levels of radioactivity associated with tail, heart, kidney, muscle and spleen remained fairly constant at 10.2% or less throughout the course of the experiment. Importantly, the radioactivity associated with the bone ranged from 4.4% of the injected dose per gram bone at 1 hour to 3.2% at 72 hours. Taken together, these results suggest that little free yttrium was associated with the conjugate and that little free radiometal was released during the course of the study. These data were used in determining radiation absorbed dose estimates for Y2B8 described below.

For tumor localization studies, 2B8-MX-DTPA was prepared and radiolabeled with  $^{111}\text{In}$  to a specific activity of 2.7 mCi/mg. One hundred microliters of labeled conjugate (approximately 24  $\mu\text{Ci}$ ) were subsequently injected into each of 12 athymic mice bearing Ramos B cell tumors. Tumors ranged in weight from 0.1 to 1.0 grams. At time points of 0, 24, 48, and 72 hours following injection, 50  $\mu\text{L}$  of blood was removed by retro-orbital puncture, the mice sacrificed by cervical dislocation, and the tail, heart, lungs, liver, kidney, spleen, muscle, femur, and tumor removed. After processing and weighing the tissues, the radioactivity associated with each tissue specimen was determined using a gamma counter and the values expressed as percent injected dose per gram.

The results (not shown) evidenced that the tumor concentrations of the  $^{111}\text{In}$ -2B8-MX-DTPA increased steadily throughout the course of the experiment. Thirteen percent of the injected dose was accumulated in the tumor after 72 hours. The blood levels, by contrast, dropped during the experiment from over 30% at time zero to 13% at 72 hours. All other tissues (except muscle) contained between 1.3 and 6.0% of the injected dose per gram tissue by the end of the experiment; muscle tissue contained approximately 13% of the injected dose per gram.

#### D. Human Studies

##### i. 2B8 and 2B8-MX-DTPA: Immunohistology Studies with Human Tissues

The tissue reactivity of murine monoclonal antibody 2B8 was evaluated using a panel of 32 different human tissues fixed with acetone. Antibody 2B8 reacts with the anti-CD20 antigen which had a very restricted pattern of tissue distribution, being observed only in a subset of cells in lymphoid tissues including those of hematopoietic origin.

In the lymph node, immunoreactivity was observed in a population of mature cortical B-lymphocytes as well as proliferating cells in the germinal centers. Positive reactivity was also observed in the peripheral blood, B-cell areas of the tonsils, white pulp of the spleen, and with 40-70% of the medullary lymphocytes found in the thymus. Positive reactivity was also seen in the follicles of the lamina propria (Peyer's Patches) of the large intestines. Finally, aggregates or scattered lymphoid cells in the stroma of various organs, including the bladder, breast, cervix, esophagus, lung, parotid, prostate, small intestine; and stomach, were also positive with antibody 2B8 (data not shown).

All simple epithelial cells, as well as the stratified epithelia and epithelia of different organs, were found to be unreactive. Similarly, no reactivity was seen with neuroectodermal cells, including those in the brain, spinal cord and peripheral nerves. Mesenchymal elements, such as skeletal and smooth muscle cells, fibroblasts, endothelial cells, and polymorphonuclear inflammatory cells were also found to be negative (data not shown).

The tissue reactivity of the 2B8-MX-DTPA conjugate was evaluated using a panel of sixteen human tissues which had been fixed with acetone. As previously demonstrated with the native antibody (data not shown), the 2B8-MX-DTPA conjugate recognized the CD20 antigen which exhibited a highly restricted pattern of distribution, being found only on a subset of cells of lymphoid origin. In the lymph node, immunoreactivity was observed in the B cell population. Strong reactivity was seen in the white pulp of the spleen and in the medullary lymphocytes of the thymus. Immunoreactivity was also observed in scattered lymphocytes in the bladder, heart, large intestines, liver, lung, and uterus, and was attributed to the presence of inflammatory cells present in these tissues. As with the native antibody, no reactivity was observed with neuroectodermal cells or with mesenchymal elements (data not shown).

##### ii. Clinical Analysis of I2B8 (Imaging) and Y2B8 (Therapy)

###### a. Phase I/II Clinical Trial Single Dose Therapy Study

A Phase I/II clinical analysis of I2B8 (imaging) followed by treatment with a single therapeutic dose of Y2B8 is currently being conducted. For the single-dose study, the following schema is being followed:

1. Peripheral Stem Cell (PSC) or Bone Marrow (BM) Harvest with Purging;
2. I2B8 Imaging;
3. Y2B8 Therapy (three Dose Levels); and
4. PSC or Autologous BM Transplantation (if necessary based upon absolute neutrophil count below  $500/\text{mm}^3$  for three consecutive days or platelets below  $20,000/\text{mm}^3$  with no evidence of marrow recovery on bone marrow examination).

The Dose Levels of Y2B8 are as follows:

Dose Level	Dose (mCi)
1.	20
2.	30
3.	40

Three patients are to be treated at each of the dose levels for determination of a Maximum Tolerated Dose ("MTD").

Imaging (Dosimetry) Studies are conducted as follows: each patient is involved in two *in vivo* biodistribution studies using I2B8. In the first study, 2mg of I2B8 (5mCi), is administered as an intravenous (i.v.) infusion over one hour; one week later 2B8 (*ie* unconjugated antibody) is administered by i.v. at a rate not to exceed 250mg/hr followed immediately by 2mg of I2B8 (5mCi) administered by i.v. over one hour. In both studies, immediately following the I2B8 infusion, each patient is imaged and imaging is repeated at time  $t = 14-18$  hr (if indicated),  $t = 24$  hr;  $t = 72$  hr; and  $t = 96$  hr (if indicated). Whole body average retention times for the indium [111] label are determined; such determinations are also made for recognizable organs or tumor lesions ("regions of interest").

The regions of interest are compared to the whole body concentrations of the label; based upon this comparison, an estimate of the localization and concentration of Y2B8 can be determined using standard protocols. If the estimated cumulative dose of Y2B8 is greater than eight (8) times the estimated whole body dose, or if the estimated cumulative dose for the liver exceeds 1500 cGy, no treatment with Y2B8 should occur.

If the imaging studies are acceptable, either 0.0 or 1.0mg/kg patient body weight of 2B8 is administered by i.v. infusion at a rate not to exceed 250mg/h. This is followed by administration of Y2B8 (10,20 or 40mCi) at an i.v. infusion rate of 20mCi/hr.

#### b. Phase I/II Clinical Trial: Multiple Dose Therapy Study

A Phase I/II clinical analysis of of Y2B8 is currently being conducted. For the multiple-dose study, the following schema is being followed:

1. PSC or BM Harvest;
2. I2B8 Imaging;
3. Y2B8 Therapy (three Dose Levels) for four doses or a total cumulative dose of 80mCi; and
4. PSC or Autologous BM Transplantation (based upon decision of medical practitioner).

The Dose Levels of Y2B8 are as follows:

Dose Level	Dose (mCi)
1.	10
2.	15
3.	20

Three patients are to be treated at each of the dose levels for determination of an MTD.

Imaging (Dosimetry) Studies are conducted as follows: A preferred imaging dose for the unlabeled antibody (*ie* 2B8) will be determined with the first two patients. The first two patients will receive 100mg of unlabeled 2B8 in 250cc of normal saline over 4 hrs followed by 0.5mCi of I2B8 --blood will be sampled for biodistribution data at times  $t = 0$ ,  $t = 10$  min.,  $t = 120$  min.,  $t = 24$  hr, and  $t = 48$  hr. Patients will be scanned with multiple regional gamma camera images at times  $t = 2$  hr,  $t = 24$  hr and  $t = 48$  hr. After scanning at  $t = 48$  hr, the patients will receive 250mg of 2B8 as described, followed by 4.5mCi of I2B8 -- blood and scanning will then follow as described. If 100mg of 2B8 produces superior imaging, then the next two patients will receive 50mg of 2B8 as described, followed by 0.5mCi of I2B8 followed 48 hrs later by 100mg 2B8 and then with 4.5mCi of I2B8. If 250mg of 2B8 produces superior imaging, then the next two patients will receive 250mg of 2B8 as described, followed by 0.5mCi of I2B8 followed 48 hrs later with 500mg 2B8 and then with 4.5mCi of I2B8. Subsequent patients will be treated with the lowest amount of 2B8 that provides optimal imaging. Optimal imaging will be defined by: (1) best effective imaging with the slowest disappearance of antibody; (2) best distribution minimizing compartmentalization in a single organ; and (3) best subjective resolution of the lesion



(tumor/background comparison).

For the first four patients, the first therapeutic dose of Y2B8 will begin 14 days after the last dose of I2B8; for subsequent patients, the first therapeutic dose of Y2B8 will begin between two to seven days after the I2B8.

Prior to treatment with Y2B8, for the patients other than the first four, 2B8 will be administered as described, followed by i.v. infusion of Y2B8 over 5-10 min. Blood will be sampled for biodistribution at times  $t = 0$ ,  $t = 10$  min.,  $t = 120$  min.,  $t = 24$  hr and  $t = 48$  hr. Patients will receive repetitive doses of Y2B8 (the same dose administered as with the first dose) approximately every six to eight weeks for a maximum of four doses, or total cumulative dose of 80mCi. It is most preferred that patients not receive a subsequent dose of Y2B8 until the patients' WBC is greater than/equal to 3,000 and AGC is greater than/equal to 100,000.

Following completion of the three-dose level study, an MTD will be defined. Additional patients will then be enrolled in the study and these will receive the MTD.

## II. CHIMERIC ANTI-CD20 ANTIBODY PRODUCTION ("C2B8")

### A. Construction of Chimeric Anti-CD20 Immunoglobulin DNA Expression Vector

RNA was isolated from the 2B8 mouse hybridoma cell (as described in Chomczynski, P. *et al.*, "Single step method of RNA isolation by acid guanidinium thiocyanate-phenol-chloroform extraction." *Anal. Biochem.* 162:156-159 (1987)). and cDNA was prepared therefrom. The mouse immunoglobulin light chain variable region DNA was isolated from the cDNA by polymerase chain reaction using a set of DNA primers with homology to mouse light chain signal sequences at the 5' end and mouse light chain J region at the 3' end. Primer sequences were as follows:

1.  $V_L$  sense (SEQ. ID. NO. 3)

5' ATC AC AGATCT CTC ACC ATG GAT TTT CAG GTG CAG ATT ATC AGC TTC 3'

(The underlined portion is a Bgl II site; the above-lined portion is the start codon.)

2.  $V_L$  Antisense (SEQ. ID. NO. 4)

5' TGC AGC ATC CGTACG TTT GAT TTC CAG CTT 3'

(The underlined portion is a Bsi WI site.)

See, Figures 1 and 2 for the corresponding Bgl II and Bsi WI sites in TCAE 8, and Figure 3 for the corresponding sites in anti-CD20 in TCAE 8.

These resulting DNA fragment was cloned directly into the TCAE 8 vector in front of the human kappa light chain constant domain and sequenced. The determined DNA sequence for the murine variable region light chain is set forth in Figure 4 (SEQ. ID. NO. 5); see also Figure 3, nucleotides 978 through 1362. Figure 4 further provides the amino acid sequence from this murine variable region, and the CDR and framework regions. The mouse light chain variable region from 2B8 is in the mouse kappa VI family. See, Kabat, *supra*.

The mouse heavy chain variable region was similarly isolated and cloned in front of the human IgG1 constant domains. Primers were as follows:

1.  $V_H$  Sense (SEQ. ID. NO. 6)

5' GCG GCT CCC ACGCGT GTC CTG TCC CAG 3'

(The underlined portion is an Mlu I site.)

2.  $V_H$  Antisense (SEQ. ID. NO. 7)

5' GG(G/C) TGT TGT GCTAGC TG(A/C) (A/G)GA GAC (G/A)GT GA 3'

(The underlined portion is an Nhe I site.)

See, Figures 1 and 2 for corresponding Mlu I and Nhe I sites in TCAE 8, and Figure 3 for corresponding sites in anti-CD20 in TCAE 8.

The sequence for this mouse heavy chain is set forth in Figure 5 (SEQ. ID. NO. 8); see also Figure 3, nucleotide 2401 through 2820. Figure 5 also provides the amino acid sequence from this murine variable region, and the CDR and framework regions. The mouse heavy chain variable region from 2B8 is in the mouse VH 2B family. See, Kabat, *supra*.

### B. Creation of Chimeric Anti-CD20 Producing CHO and SP2/0 Transfectomas

Chinese hamster ovary ("CHO") cells DG44 were grown in SSFM II minus hypoxanthine and thymidine media (Gibco, Grand Islands, NY, Form No. 91-0456PK); SP2/0 mouse myeloma cells were grown in Dulbecco's Modified Eagles Medium media ("DMEM") (Irvine Scientific, Santa Ana, Ca., Cat. No. 9024) with 5% fetal bovine serum and 20

ml/L glutamine added. Four million cells were electroporated with either 25 µg CHO or 50 µg SP2/0 plasmid DNA that had been restricted with Not I using a BTX 600 electroporation system (BTX, San Diego, CA) in 0.4 ml disposable cuvettes. Conditions were either 210 volts for CHO or 180 volts for SP2/0, 400 microfaradays, 13 ohms. Each electroporation was plated into six 96 well dishes (about 7,000 cells/well). Dishes were fed with media containing G418 (GENETICIN, Gibco, Cat. No. 860-1811) at 400 µg/ml active compound for CHO (media further included 50 µM hypoxanthine and 8 µM thymidine) or 800 µg/ml for SP2/0, two days following electroporation and thereafter 2 or 3 days until colonies arose. Supernatant from colonies was assayed for the presence of chimeric immunoglobulin via an ELISA specific for human antibody. Colonies producing the highest amount of immunoglobulin were expanded and plated into 96 well plates containing media plus methotrexate (25 nM for SP2/0 and 5nM for CHO) and fed every two or three days. Supernatants were assayed as above and colonies producing the highest amount of immunoglobulin were examined. Chimeric anti-CD20 antibody was purified from supernatant using protein A affinity chromatography.

Purified chimeric anti-CD20 was analyzed by electrophoresis in polyacrylamide gels and estimated to be greater than about 95% pure. Affinity and specificity of the chimeric antibody was determined based upon 2B8. Chimeric anti-CD20 antibody tested in direct and competitive binding assays, when compared to murine anti-CD20 monoclonal antibody 2B8, evidenced comparable affinity and specificity on a number of CD20 positive B cells lines (data not presented). The apparent affinity constant ("Kap") of the chimeric antibody was determined by direct binding of  $^{125}$ I radiolabeled chimeric anti-CD20 and compared to radiolabeled 2B8 by Scatchard-plot; estimated Kap for CHO produced chimeric anti-CD20 was  $5.2 \times 10^{-9}$  M and for SP2/0 produced antibody,  $7.4 \times 10^{-9}$  M. The estimated Kap for 2B8 was  $3.5 \times 10^{-9}$  M. Direct competition by radioimmunoassay was utilized to confirm both the specificity and retention of immunoreactivity of the chimeric antibody by comparing its ability to effectively compete with 2B8. Substantially equivalent amounts of chimeric anti-CD20 and 2B8 antibodies were required to produce 50% inhibition of binding to CD20 antigens on B cells (data not presented), *ie* there was a minimal loss of inhibiting activity of the anti-CD20 antibodies, presumably due to chimerization.

The results of Example II.B indicate, *inter alia*, that chimeric anti-CD20 antibodies were generated from CHO and SP2/0 transfectomas using the TCAE 8 vectors, and these chimeric antibodies had substantially the same specificity and binding capability as murine anti-CD20 monoclonal antibody 2B8.

### C. Determination of Immunological Activity of Chimeric Anti-CD20 Antibodies

#### i. Human C1q Analysis

Chimeric anti-CD20 antibodies produced by both CHO and SP2/0 cell lines were evaluated for human C1q binding in a flow cytometry assay using fluorescein labeled C1q (C1q was obtained from Quidel, Mira Mesa, CA, Prod. No. A400 and FITC label from Sigma, St. Louis MO, Prod. No. F-7250; FITC. Labeling of C1q was accomplished in accordance with the protocol described in *Selected Methods In Cellular Immunology*, Mitchell & Shiigi, Ed. (W.H. Freeman & Co., San Francisco, CA, 1980, p. 292). Analytical results were derived using a Becton Dickinson FACScan™ flow cytometer (fluorescein measured over a range of 515-545 nm). Equivalent amounts of chimeric anti-CD20 antibody, human IgG1,K myeloma protein (Binding Site, San Diego, Ca, Prod. No. BP078), and 2B8 were incubated with an equivalent number of CD20-positive SB cells, followed by a wash step with FACS buffer (.2% BSA in PBS, pH 7.4, .02% sodium azide) to remove unattached antibody, followed by incubation with FITC labeled C1q. Following a 30-60 min. incubation, cells were again washed. The three conditions, including FITC-labeled C1q as a control, were analyzed on the FACScan™ following manufacturing instructions. Results are presented in Figure 6.

As the results of Figure 6 evidence, a significant increase in fluorescence was observed only for the chimeric anti-CD20 antibody condition; *ie* only SB cells with adherent chimeric anti-CD20 antibody were C1q positive, while the other conditions produced the same pattern as the control.

#### ii. Complement Dependent Cell Lyses

Chimeric anti-CD20 antibodies were analyzed for their ability to lyse lymphoma cell lines in the presence of human serum (complement source). CD20 positive SB cells were labeled with  $^{51}$ Cr by admixing 100µ Ci of  $^{51}$ Cr with  $1 \times 10^6$  SB cells for 1 hr at 37°C; labeled SB cells were then incubated in the presence of equivalent amounts of human complement and equivalent amounts (0-50 µg/ml) of either chimeric anti-CD20 antibodies or 2B8 for 4 hrs at 37°C (see, Brunner, K.T. *et al.*, "Quantitative assay of the lytic action of immune lymphoid cells on  $^{51}$ Cr-labeled allogeneic target cells *in vitro*." *Immunology* 14:181-189 (1968). Results are presented in Figure 7.

The results of Figure 7 indicate, *inter alia*, that chimeric anti-CD20 antibodies produced significant lysis (49%) under these conditions.

### iii. Antibody Dependent Cellular Cytotoxicity Effector Assay

For this study, CD20 positive cells (SB) and CD20 negative cells (T cell leukemia line HSB; *see*, Adams, Richard, "Formal Discussion," *Can. Res.* 27:2479-2482 (1967); ATCC deposit no. ATCC CCL 120.1) were utilized; both were labeled with  $^{51}\text{Cr}$ . Analysis was conducted following the protocol described in Brunner, K.T. *et al.*, "Quantitative assay of the lytic action of immune lymphoid cells on  $^{51}\text{Cr}$ -labeled allogeneic target cells *in vitro*; inhibition by isoantibody and drugs." *Immunology* 14:181-189 (1968); a substantial chimeric anti-CD20 antibody dependent cell mediated lysis of CD20 positive SB target cells ( $^{51}\text{Cr}$ -labeled) at the end of a 4 hr, 37°C incubation, was observed and this effect was observed for both CHO and SP2/0 produced antibody (effector cells were human peripheral lymphocytes; ratio of effector cells:target was 100:1). Efficient lysis of target cells was obtained at 3.9 µg/ml. In contrast, under the same conditions, the murine anti-CD20 monoclonal antibody 2B8 had a statistically insignificant effect, and CD20 negative HSB cells were not lysed. Results are presented in Figure 8.

The results of Example II indicate, *inter alia*, that the chimeric anti-CD20 antibodies of Example I were immunologically active.

### III. DEPLETION OF B CELLS *IN VIVO* USING CHIMERIC ANTI-CD20

#### A. Non-Human Primate Study

Three separate non-human primate studies were conducted. For convenience, these are referred to herein as "Chimeric Anti-CD20: CHO & SP2/0," "Chimeric Anti-CD20: CHO," and "High Dosage Chimeric Anti-CD20." Conditions were as follows:

#### Chimeric Anti-CD20: CHO & SP2/0

Six cynomolgus monkeys ranging in weight from 4.5 to 7 kilograms (White Sands Research Center, Alamogordo, NM) were divided into three groups of two monkeys each. Both animals of each group received the same dose of immunologically active chimeric anti-CD20 antibody. One animal in each group received purified antibody produced by the CHO transfectoma; the other received antibody produced by the SP2/0 transfectoma. The three groups received antibody dosages corresponding to 0.1 mg/kg, 0.4 mg/kg, and 1.6 mg/kg each day for four (4) consecutive days. The chimeric immunologically active anti-CD20 antibody, which was admixed with sterile saline, was administered by intravenous infusion; blood samples were drawn prior to each infusion. Additional blood samples were drawn beginning 24 hrs after the last injection (T=0) and thereafter on days 1, 3, 7, 14 and 28; blood samples were also taken thereafter at biweekly intervals until completion of the study at day 90.

Approximately 5 ml of whole blood from each animal was centrifuged at 2000 RPM for 5 min. Plasma was removed for assay of soluble chimeric anti-CD20 antibody levels. The pellet (containing peripheral blood leukocytes and red blood cells) was resuspended in fetal calf serum for fluorescent-labeled antibody analysis (*see*, "Fluorescent Antibody Labeling of Lymphoid Cell Population," *infra.*).

#### Chimeric Anti-CD20: CHO

Six cynomolgus monkeys ranging in weight from 4 to 6 kilograms (White Sands) were divided into three groups of two monkeys each. All animals were injected with immunologically active chimeric anti-CD20 antibodies produced from the CHO transfectoma (in sterile saline). The three groups were separated as follows: subgroup 1 received daily intravenous injections of 0.01 mg/kg of the antibody over a four (4) day period; subgroup 2 received daily intravenous injections of 0.4 mg/kg of the antibody over a four (4) day period; subgroup 3 received a single intravenous injection of 6.4 mg/kg of the antibody. For all three subgroups, a blood sample was obtained prior to initiation of treatment; additionally, blood samples were also drawn at T=0, 1, 3, 7, 14 and 28 days following the last injection as described above, and these samples were processed for fluorescent labeled antibody analysis (*see*, "Fluorescent Antibody Labeling," *infra.*). In addition to peripheral blood B cell quantitation, lymph node biopsies were taken at days 7, 14 and 28 following the last injection, and a single cell preparation stained for quantitation of lymphocyte populations by flow cytometry.

#### High Dosage Chimeric Anti-CD20

Two cynomolgus monkeys (White Sands) were infused with 16.8 mg/kg of the immunologically active chimeric anti-CD20 antibodies from the CHO transfectomas (in sterile saline) weekly over a period of four consecutive weeks. At the conclusion of the treatment, both animals were anesthetized for removal of bone marrow; lymph node biopsies were also taken. Both sets of tissue were stained for the presence of B lymphocytes using Leu 16 by flow cytometry following the protocol described in Ling, N.R. *et al.*, "B-cell and plasma cell antigens." *Leucocyte Typing III White Cell Differenti-*

ations Antigens, A.J. McMichael, Ed. (Oxford University Press, Oxford UK, 1987), p. 302.

#### Fluorescent Antibody Labeling of Lymphoid Cell Population

After removal of plasma, leukocytes were washed twice with Hanks Balanced Salt Solution ("HBSS") and resuspended in a plasma equivalent volume of fetal bovine serum (heat inactivated at 56°C for 30 min.). A 0.1 ml volume of the cell preparation was distributed to each of six (6), 15 ml conical centrifuge tubes. Fluorescein labeled monoclonal antibodies with specificity for the human lymphocyte surface markers CD2 (AMAC, Westbrook, ME), CD20 (Becton Dickinson) and human IgM (Binding Site, San Diego, CA) were added to 3 of the tubes for identifying T and B lymphocyte populations. All reagents had previously tested positive to the corresponding monkey lymphocyte antigens. Chimeric anti-CD20 antibody bound to monkey B cell surface CD20 was measured in the fourth tube using polyclonal goat anti-human IgG coupled with phycoerythrin (AMAC). This reagent was pre-adsorbed on a monkey Ig-sepharose column to prevent cross-reactivity to monkey Ig, thus allowing specific detection and quantitation of chimeric anti-CD20 antibody bound to cells. A fifth tube included both anti-IgM and anti-human IgG reagents for double stained B cell population. A sixth sample was included with no reagents for determination of autofluorescence. Cells were incubated with fluorescent antibodies for 30 min., washed and fixed with 0.5 ml of fixation buffer (0.15 M NaCl, 1% paraformaldehyde, pH7.4) and analyzed on a Becton Dickinson FACScan™ instrument. Lymphocyte populations were initially identified by forward versus right angle light scatter in a dot-plot bitmap with unlabeled leucocytes. The total lymphocyte population was then isolated by gating out all other events. Subsequent fluorescence measurements reflected only gated lymphocyte specific events.

#### Depletion of Peripheral Blood B Lymphocytes

No observable difference could be ascertained between the efficacy of CHO and SP2/0 produced antibodies in depleting B cells *in vivo*, although a slight increase in B cell recovery beginning after day 7 for monkeys injected with chimeric anti-CD20 antibodies derived from CHO transfectomas at dosage levels 1.6 mg/kg and 6.4 mg/kg was observed and for the monkey injected with SP2/0 producing antibody at the 0.4 mg/kg dose level. Figures 9A, B and C provide the results derived from the chimeric anti-CD20:CHO & SP2/0 study, with Figure 9A directed to the 0.4 mg/kg dose level; Figure 9B directed to the 1.6 mg/kg dose level; and Figure 9C directed to the 6.4 mg/kg dose level.

As is evident from Figure 9, there was a dramatic decrease (>95%) in peripheral B cell levels after the therapeutic treatment across all tested dose ranges, and these levels were maintained up to seven (7) days post infusion; after this period, B cell recovery began, and, the time of recovery initiation was independent of dosage levels.

In the Chimeric Anti-CD20:CHO study, a 10-fold lower antibody dosage concentration (0.01 mg/kg) over a period of four daily injections (0.04 mg/kg total) was utilized. Figure 10 provides the results of this study. This dosage depleted the peripheral blood B cell population to approximately 50% of normal levels estimated with either the anti-surface IgM or the Leu 16 antibody. The results also indicate that saturation of the CD20 antigen on the B lymphocyte population was not achieved with immunologically active chimeric anti-CD20 antibody at this dose concentration over this period of time for non-human primates; B lymphocytes coated with the antibody were detected in the blood samples during the initial three days following therapeutic treatment. However, by day 7, antibody coated cells were undetectable.

Table I summarizes the results of single and multiple doses of immunologically active chimeric anti-CD20 antibody on the peripheral blood populations; single dose condition was 6.4 mg/kg; multiple dose condition was 0.4 mg/kg over four (4) consecutive days (these results were derived from the monkeys described above).

TABLE I

PERIPHERAL BLOOD POPULATION FROM C2B8 PRIMATE STUDY				
Monkey	Dose	Day	CD2	Anti-Hu IgG
A	0.4 mg/kg (4 doses)	Prebleed	81.5	-
		0	86.5	0.2
		7	85.5	0.0
		21	93.3	-
		28	85.5	-

TABLE I (continued)

PERIPHERAL BLOOD POPULATION FROM C2B8 PRIMATE STUDY				
Monkey	Dose	Day	CD2	Anti-Hu IgG
B	0.4 mg/kg (4 doses)	Prebleed	81.7	-
		0	94.6	0.1
		7	92.2	0.1
		21	84.9	-
		28	84.1	-
C	6.4 mg/kg (1 dose)	Prebleed	77.7	0.0
		7	85.7	0.1
		21	86.7	-
		28	76.7	-
D	6.4 mg/kg (1 dose)	Prebleed	85.7	0.1
		7	94.7	0.1
		21	85.2	-
		28	85.9	-
Monkey	Anti-Hu IgG+ Anti-Hu IgM*	Leu-16	% B Cell Depletion	
A	-	9.4	0	
	0.3	0.0	97	
	0.1	1.2	99	
	-	2.1	78	
	-	4.1	66	
B	-	14.8	0	
	0.2	0.1	99	
	0.1	0.1	99	
	-	6.9	53	
	-	8.7	41	
C	0.2	17.0	0	
	0.1	0.0	99	
	-	14.7	15	
	-	8.1	62	
D	0.1	14.4	0	
	0.2	0.0	99	
	-	9.2	46	
	-	6.7	53	

\*Double staining population which indicates extent of chimeric anti-CD20 coated B cells.

The data summarized in Table I indicates that depletion of B cells in peripheral blood under conditions of antibody excess occurred rapidly and effectively, regardless of single or multiple dosage levels. Additionally, depletion was observed for at least seven (7) days following the last injection, with partial B cell recovery observed by day 21.

Table II summarizes the effect of immunologically active, chimeric anti-CD20 antibodies on cell populations of

# EP 0 752 248 A1

lymph nodes using the treatment regimen of Table I (4 daily doses of 0.4 mg/kg; 1 dose of 6.4 mg/kg); comparative values for normal lymph nodes (control monkey, axillary and inguinal) and normal bone marrow (two monkeys) are also provided.

TALE II

CELL POPULATIONS OF LYMPH NODES						
Monkey	Dose		Day	CD2	Anti-Hu IgM	
A	0.4 mg/kg (4 doses)		7	66.9	-	
			14	76.9	19.6	
			28	61.6	19.7	
B	0.4 mg/kg (4 doses)		7	59.4	-	
			14	83.2	9.9	
			28	84.1	15.7	
C	6.4 mg/kg (1 dose)		7	75.5	-	
			14	74.1	17.9	
			28	66.9	23.1	
D	6.4 mg/kg (1 dose)		7	83.8	-	
			14	74.1	17.9	
			28	84.1	12.8	
Monkey	Anti-Hu IgG + Anti-Hu IgM		Leu-16	% B Lymphocyte Depletion		
A	7.4		40.1	1		
	0.8		22.6	44		
	-		26.0	36		
B	29.9		52.2	0		
	0.7		14.5	64		
	-		14.6	64		
C	22.3		35.2	13		
	1.1		23.9	41		
	-		21.4	47		
D	12.5		19.7	51		
	0.2		8.7	78		
	-		12.9	68		
	CD2	Anti-Hu IgG+ Anti-Hu IgM		Anti-Hu IgM	Leu-16	% B Lymphocyte Depletion
Normal Lymph Nodes						
Control 1						
Axillary	55.4	25.0		-	41.4	NA
Inguinal	52.1	31.2		-	39.5	NA
Normal Bone Marrow						
Control 2	65.3	19.0		-	11.4	NA
Control 3	29.8	28.0		-	16.6	NA

The results of Table II evidence effective depletion of B lymphocytes for both treatment regimens. Table II further

indicates that for the non-human primates, complete saturation of the B cells in the lymphatic tissue with immunologically active, chimeric anti-CD20 antibody was not achieved; additionally, antibody coated cells were observed seven (7) days after treatment, followed by a marked depletion of lymph node B cells, observed on day 14.

Based upon this data, the single High Dosage Chimeric Anti-CD20 study referenced above was conducted, principally with an eye toward pharmacology/toxicology determination. *ie* this study was conducted to evaluate any toxicity associated with the administration of the chimeric antibody, as well as the efficacy of B cell depletion from peripheral blood lymph nodes and bone marrow. Additionally, because the data of Table II indicates that for that study, the majority of lymph node B cells were depleted between 7 and 14 days following treatment, a weekly dosing regimen might evidence more efficacious results. Table III summarizes the results of the High Dosage Chimeric Anti-CD20 study.

TABLE III

CELL POPULATIONS OF LYMPH NODES AND BONE MARROW					
Lymphocyte Populations (%)					
Monkey	CD2	CD20 <sup>a</sup>	mIgM + anti-C2B8 <sup>b</sup>	C2B8 <sup>c</sup>	Day <sup>d</sup>
<i>Inguinal Lymph Node</i>					
E	90.0	5.3	4.8	6.5	22
F	91.0	6.3	5.6	6.3	22
G	89.9	5.0	3.7	5.8	36
H	85.4	12.3	1.7	1.8	36
<i>Bone Marrow</i>					
E	46.7	4.3	2.6	2.8	22
F	41.8	3.0	2.1	2.2	22
G	35.3	0.8	1.4	1.4	36
H	25.6	4.4	4.3	4.4	36

<sup>a</sup>Indicates population stained with Leu 16.

<sup>b</sup>Indicates double staining population, positive for surface IgM cells and chimeric antibody coated cells.

<sup>c</sup>Indicates total population staining for chimeric antibody including double staining surface IgM positive cells and single staining (surface IgM negative) cells.

<sup>d</sup>Days after injection of final 16.8 mg/kg dose.

Both animals evaluated at 22 days post treatment cessation contained less than 5% B cells, as compared to 40% in control lymph nodes (*see*, Table II, *supra*). Similarly, in the bone marrow of animals treated with chimeric anti-CD20 antibody, the levels of CD20 positive cells were less than 3% as compared to 11-15% in the normal animals (*see*, Table II, *supra*). In the animals evaluated at 36 days post treatment cessation, one of the animals (H) had approximately 12% B cells in the lymph node and 4.4% B cells in bone marrow, while the other (G) had approximately 5% B cells in the lymph node and 0.8% in the bone marrow--the data is indicative of significant B cell depletion.

The results of Example III.A indicate, *inter alia*, that low doses of immunologically active, chimeric anti-CD20 leads to long-term peripheral blood B cell depletion in primates. The data also indicates that significant depletion of B cell populations was achieved in peripheral lymph nodes and bone marrow when repetitive high doses of the antibody were administered. Continued follow-up on the test animals has indicated that even with such severe depletion of peripheral B lymphocytes during the first week of treatment, no adverse health effects have been observed. Furthermore, as recovery of B cell population was observed, a conclusion to be drawn is that the pluripotent stem cells of these primates were not adversely affected by the treatment.

#### B. Clinical Analysis of C2B8

##### i. Phase I/II Clinical Trial of C2B8: Single Dose Therapy Study

Fifteen patients having histologically documented relapsed B cell lymphoma have been treated with C2B8 in a Phase I/II Clinical Trial. Each patient received a single dose of C2B8 in a dose-escalating study; there were three patients per dose: 10mg/m<sup>2</sup>; 50mg/m<sup>2</sup>; 100mg/m<sup>2</sup>; 250mg/m<sup>2</sup> and 500mg/m<sup>2</sup>. Treatment was by i.v. infusion through

an 0.22 micron in-line filter with C2B8 being diluted in a final volume of 250cc or a maximal concentration of 1mg/ml of normal saline. Initial rate was 50cc/hr for the first hour; if no toxicity was seen, dose rate was able to be escalated to a maximum of 200cc/hr.

Toxicity (as indicated by the clinician) ranged from "none", to "fever" to "moderate" (two patients) to "severe" (one patient); all patients completed the therapy treatment. Peripheral Blood Lymphocytes were analyzed to determine, *inter alia*, the impact of C2B8 on T-cells and B-cells. Consistently for all patients: Peripheral Blood B Lymphocytes were depleted after infusion with C2B8 and such depletion was maintained for in excess of two weeks.

One patient (receiving 100mg/m<sup>2</sup> of C2B8) evidenced a Partial Response to the C2B8 treatment (reduction of greater than 50% in the sum of the products of the perpendicular diameters of all measurable indicator lesions lasting greater than four weeks, during which no new lesions may appear and no existing lesions may enlarge); at least one other patient (receiving 500mg/m<sup>2</sup>) evidenced a Minor Response to the C2B8 treatment (reduction of less than 50% but at least 25% in the sum of the products of the two longest perpendicular diameters of all measurable indicator lesions). For presentational efficiency, results of the PBLs are set forth in Figure 14; data for the patient evidencing a PR is set forth in Figure 14A; for the patient evidencing an MR, data is set forth in Figure 14B. In Figure 14, the following are applicable: ■ = Lymphocytes; □ = CD3+ cells (T cells); ▲ = CD20+ cells; ● = CD19+ cells; ⊖ = Kappa; △ = lambda; and ◆ = C2B8. As evidenced, the B cell markers CD20 and CD19, Kappa and Lambda, were depleted for a period in excess of two weeks; while there was a slight, initial reduction in T-cell counts, these returned to an approximate baseline level in a relatively rapid time-frame.

#### ii. Phase I/II Clinical Trial of C2B8: Multiple Dose Therapy Study

Patients having histologically confirmed B cell lymphoma with measurable progressive disease are eligible for this study which is separated into two parts: in Phase I, consisting of a dose escalation to characterize dose limiting toxicities and determination of biologically active tolerated dose level, groups of three patients will receive weekly i.v. infusions of C2B8 for a total of four (4) separate infusions. Cumulative dose at each of the three levels will be as follows: 500mg/m<sup>2</sup> (125mg/m<sup>2</sup>/infusion); 1000mg/m<sup>2</sup> (250mg/m<sup>2</sup>/infusion); 1500mg/m<sup>2</sup> (375mg/m<sup>2</sup>/infusion. A biologically active tolerated dose is defined, and will be determined, as the lowest dose with both tolerable toxicity and adequate activity); in Phase II, additional patients will receive the biologically active tolerated dose with an emphasis on determining the activity of the four doses of C2B8.

#### IV. COMBINATION THERAPY: C2B8 AND Y2B8

A combination therapeutic approach using C2B8 and Y2B8 was investigated in a mouse xenographic model (nu/nu mice, female, approximately 10 weeks old) utilizing a B cell lymphoblastic tumor (Ramos tumor cells). For comparative purposes, additional mice were also treated with C2B8 and Y2B8.

Ramos tumor cells (ATCC, CRL 1596) were maintained in culture using RPMI-1640 supplemented with 10% fetal calf serum and glutamine at 37°C and 5% CO<sub>2</sub>. Tumors were initiated in nine female nude mice approximately 7-10 weeks old by subcutaneous injection of 1.7 x 10<sup>6</sup> Ramos cells in a volume of 0.10ml (HBSS) using a 1cc syringe fitted with 25g needle. All animals were manipulated in a laminar flow hood and all cages, bedding, food and water were autoclaved. Tumor cells were passaged by excising tumors and passing these through a 40 mesh screen; cells were washed twice with 1X HBSS (50ml) by centrifugation (1300RPM), resuspended in 1X HBSS to 10 x 10<sup>6</sup> cells/ml, and frozen at -70°C until used.

For the experimental conditions, cells from several frozen lots were thawed, pelleted by centrifugation (1300RPM) and washed twice with 1X HBSS. Cells were then resuspended to approximately 2.0 x 10<sup>6</sup> cells/ml. Approximately 9 to 12 mice were injected with 0.10ml of the cell suspension (s.c.) using a 1cc syringe fitted with a 25g needle; injections were made on the animal's left side, approximately mid-region. Tumors developed in approximately two weeks. Tumors were excised and processed as described above. Study mice were injected as described above with 1.67 x 10<sup>6</sup> cells in 0.10ml HBSS.

Based on preliminary dosing experiments, it was determined that 200mg of C2B8 and 100μCi of Y2B8 would be utilized for the study. Ninety female nu/nu mice (approximately 10 weeks old) were injected with the tumor cells. Approximately ten days later, 24 mice were assigned to four study groups (six mice/group) while attempting to maintain a comparable tumor size distribution in each group (average tumor size, expressed as a product of length x width of the tumor, was approximately 80mm<sup>2</sup>). The following groups were treated as indicated via tail-vein injections using a 100μl Hamilton syringe fitted with a 25g needle:

- A. Normal Saline
- B. Y2B8 (100μCi)
- C. C2B8 (200μg); and
- D. Y2B8 (100μCi) + C2B8 (200μg)



Groups tested with C2B8 were given a second C2B8 injection (200 $\mu$ g/mouse) seven days after the initial injection. Tumor measurements were made every two or three days using a caliper.

Preparation of treatment materials were in accordance with the following protocols:

#### 5 A. Preparation of Y2B8

Yttrium-[90] chloride (6mCi) was transformed to a polypropylene tube and adjusted to pH 4.1-4.4 using metal free 2M sodium acetate. 2B8-MX-DTPA (0.3mg in normal saline; *see above* for preparation of 2B8-MX-DTPA) was added and gently mixed by vortexing. After 15 min. incubation, the reaction was quenched by adding 0.05 x volume 20mM EDTA and 0.05X volume 2M sodium acetate. Radioactivity concentration was determined by diluting 5.0 $\mu$ l of the reaction mixture in 2.5ml 1 x PBS containing 75mg/ml HSA and 1mM DTPA ("formulation buffer"); counting was accomplished by adding 10.0 $\mu$ l to 20ml of Ecolume™ scintillation cocktail. The remainder of the reactive mixture was added to 3.0ml formulation buffer, sterile filtered and stored at 2-8°C until used. Specific activity (14mCi/mg at time of injection) was calculated using the radioactivity concentration and the calculated protein concentration based upon the amount of antibody added to the reaction mixture. Protein-associated radioactivity was determined using instant thin-layer chromatography. Radioincorporation was 95%. Y2B8 was diluted in formulation buffer immediately before use and sterile-filtered (final radioactivity concentration was 1.0mCi/ml).

#### 20 B. Preparation of C2B8

C2B8 was prepared as described above. C2B8 was provided as a sterile reagent in normal saline at 5.0mg/ml. Prior to injection; the C2B8 was diluted in normal saline to 2.0mg/ml and sterile filtered.

#### 25 C. Results

Following treatment, tumor size was expressed as a product of length and width, and measurements were taken on the days indicated in Figure 11 (Y2B8 vs. Saline); Figure 12 (C2B8 vs. Saline); and Figure 13 (Y2B8 + C2B8 vs. Saline). Standard error was also determined.

As indicated in Figure 13, the combination of Y2B8 and C2B8 exhibited tumoricidal effects comparable to the effects evidenced by either Y2B8 or C2B8.

### V. ALTERNATIVE THERAPY STRATEGIES

Alternative therapeutic strategies recognized in view of the foregoing examples are evident. One such strategy employs the use of a therapeutic dose of C2B8 followed within about one week with a combination of either 2B8 and radioabeled 2B8 (*eg* Y2B8); or 2B8, C2B8 and, *eg* Y2B8; or C2B8 and, *eg* Y2B8. An additional strategy is utilization of radiolabeled C2B8 -- such a strategy allows for utilization of the benefits of the immunologically active portion of C2B8 plus those benefits associated with a radiolabel. Preferred radiolabels include yttrium-90 given the larger circulating half-life of C2B8 versus the murine antibody 2B8. Because of the ability of C2B8 to deplete B-cells, and the benefits to be derived from the use of a radiolabel, a preferred alternative strategy is to treat the patient with C2B8 (either with a single dose or multiple doses) such that most, if not all, peripheral B cells have been depleted. This would then be followed with the use of radiolabeled 2B8; because of the depletion of peripheral B cells, the radiolabeled 2B8 stands an increased chance of targeting tumor cells. Iodine [131] labeled 2B8 is preferably utilized, given the types of results reported in the literature with this label (*see* Kaminski). An alternative preference involves the use of a radiolabeled 2B8 (or C2B8) first in an effort to increase the permeability of a tumor, followed by single or multiple treatments with C2B8; the intent of this strategy is to increase the chances of the C2B8 in getting both outside and inside the tumor mass. A further strategy involved the use of chemotherapeutic agent in combination with C2B8. These strategies include so-called "staggered" treatments, *ie*, treatment with chemotherapeutic agent, followed by treatment with C2B8, followed by a repetition of this protocol. Alternatively, initial treatment with a single or multiple doses of C2B8, thereafter followed with chemotherapeutic treatment, is viable. Preferred chemotherapeutic agents include, but are not limited to: cyclophosphamide; doxorubicin; vincristine; and prednisone, *See* Armitage, J.O. *et al*, *Cancer* 50:1695 (1982), incorporated herein by reference.

The foregoing alternative therapy strategies are not intended to be limiting, but rather are presented as being representative.

### 55 VI. DEPOSIT INFORMATION

Anti-CD20 in TCAE 8 (transformed in *E. coli* for purposes of deposit) was deposited with the American Type Culture Collection (ATCC), 12301 Parklawn Drive, Rockville, Maryland, 20852, under the provisions of the Budapest Treaty for

the International Recognition of the Deposit of Microorganisms for the Purpose of Patent Procedure ("Budapest Treaty"). The microorganism was tested by the ATCC on November 9, 1992, and determined to be viable on that date. The ATCC has assigned this microorganism for the following ATCC deposit number: ATCC 69119 (anti-CD20 in TCAE 8). Hybridoma 2B8 was deposited with the ATCC on June 22, 1993 under the provisions of the Budapest Treaty. The  
 5 viability of the culture was determined on June 25, 1993 and the ATCC has assigned this hybridoma the following ATCC deposit number: HB 11388.

## G. SEQUENCE LISTING

### (1) GENERAL INFORMATION

(i) APPLICANT: Darrell Anderson, Nabil Hanna, John Leonard,  
 15 Roland Newman and Mitchell Reff and William H. Rastetter

(ii) TITLE OF INVENTION: THERAPEUTIC APPLICATION OF  
 20 CHIMERIC AND RADIOLABELED  
 ANTIBODIES TO HUMAN B  
 LYMPHOCYTE RESTRICTED  
 DIFFERENTIATION ANTIGEN FOR  
 TREATMENT OF B CELL LYMPHOMA

(iii) NUMBER OF SEQUENCES: 8

(iv) CORRESPONDING ADDRESS:

(A) ADDRESSEE: IDEC Pharmaceuticals Corporation  
 30 (B) STREET: 11011 Torreyana Road  
 (C) CITY: San Diego  
 (D) STATE: California  
 (E) COUNTRY: USA  
 (F) ZIP: 92121

(v) COMPUTER READABLE FORM:

(A) MEDIUM TYPE: Diskette, 3.5 inch, 1.44 Mb  
 40 (B) COMPUTER: Macintosh  
 (C) OPERATING SYSTEM: MS.DOS  
 (D) SOFTWARE: Microsoft Word 5.0

(vi) CURRENT APPLICATION DATA:

(A) APPLICATION NUMBER:  
 45 (B) FILING DATE:  
 (C) CLASSIFICATION:

(viii) ATTORNEY/AGENT INFORMATION:

(A) NAME: Burgoon, Richard P. Jr.  
 50 (B) REGISTRATION NUMBER: 34,787  
 (C) REFERENCE/DOCKET NUMBER:

(ix) TELECOMMUNICATION INFORMATION:

(A) TELEPHONE: (619) 550-8500  
 55 (B) TELEFAX: (619) 550-8750

## (2) INFORMATION FOR SEQ ID NO: 1:

## (i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 8540 bases  
 (B) TYPE: nucleic acid  
 (C) STRANDEDNESS: single  
 (D) TOPOLOGY: circular

## (ii) MOLECULE TYPE: DNA (genomic)

## (iii) HYPOTHETICAL: yes

## (iv) ANTI-SENSE: no

## (ix) SEQUENCE DESCRIPTION: SEQ ID NO: 1:

```

GACGTCGCGG CCGCTCTAGG CCTCCAAAA AGCCTCCTCA CTACTTCTGG AATAGCTCAG      60
AGGCCGAGGC GGCCTCGGCC TCTGCATAAA TAAAAAAAT TAGTCAGCCA TGCATGGGGC      120
GGAGAATGGG CGGAAGTGGG CGGAGTTAGG GGCGGGATGG GCGGAGTTAG GGGCGGGACT      180
ATGTTTGCTG ACTAATTGAG ATGCATGCTT TGCATACTTC TGCCTGCTGG GGAGCCTGGG      240
GACTTTCCAC ACCTGGTTGC TGACTAATTG AGATGCATGC TTTGCATACT TCTGCCTGCT      300
GGGGAGCCTG GGGACTTTCC ACACCCTAAC TGACACACAT TCCACAGAAT TAATTCCCCT      360
AGTTATTAAT AGTAATCAAT TACGGGGTCA TTAGTTCATA GCCCATATAT GGAGTTCGCG      420
GTTACATAAC TTACGGTAAA TGGCCCGCCT GGCTGACCGC CCAACGACCC CCGCCCATG      480
ACGTCAATAA TGACGTATGT TCCCATAGTA ACGCCAATAG GGACTTTCCA TTGACGTCAA      540
TGGGTGGACT ATTTACGGTA AACTGCCCAC TTGGCAGTAC ATCAAGTGTA TCATATGCCA      600
AGTACGCCCC CTATTGACGT CAATGACGGT AAATGGCCCC CCTGGCATTG TGCCAGTAC      660
ATGACCTTAT GGGACTTTCC TACTTGGCAG TACATCTACG TATTAGTCAT CGCTATTACC      720
ATGGTGATGC GGTTTTGGCA GTACATCAAT GGGCGTGGAT AGCGGTTTGA CTCACGGGGA      780
TTTCCAAGTC TCCACCCCAT TGACGTCAAT GGGAGTTTGT TTTGGCACCA AAATCAACGG      840
GACTTTCCAA AATGTCGTAA CAACTCCGCC CCATTGACGC AAATGGGCGG TAGGCGTGTA      900
CGGTGGGAGG TCTATATAAG CAGAGCTGGG TACGTGAACC GTCAGATCGC CTGGAGACGC      960
CATCACAGAT CTCTACCAT GAGGGTCCCC GCTCAGCTCC TGGGGCTCCT GCTGCTCTGG     1020
CTCCAGGTG CACGATGTGA TGGTACCAAG GTGGAATCA AACGTACGGT GGCTGCACCA     1080
TCTGTCTTCA TCTTCCCGCC ATCTGATGAG CAGTTGAAAT CTGGAACTGC CTCTGTGTGT     1140
TGCCTGCTGA ATAATTCTA TCCCAGAGAG GCCAAAGTAC AGTGAAGGT GGATAACGCC     1200
CTCCAATCGG GTAATCCCA GGAGAGTGTC ACAGAGCAGG ACAGCAAGGA CAGCACCTAC     1260

```

EP 0 752 248 A1

AGCCTCAGCA GCACCCCTGAC GCTGAGCAAA GCAGACTACG AGAAACACAA AGTCTACGCC 1332  
 TGCGAAGTCA CCCATCAGGG CCTGAGCTCG CCCGTCACAA AGAGCTTCAA CAGGGGAGAG 1380  
 5 TGTGTAATTC AGATCCGTTA ACGGTTACCA ACTACCTAGA CTGGATTTCGT GACAACATGC 1440  
 GGCCGTGATA TCTACGTATG ATCAGCCTCG ACTGTGCTTT CTAGTTGCCA GCCATCTGTT 1500  
 GTTTGCCCTT CCCCCGTGCC TTCCTTGACC CTGGAAGGTG CCACTCCAC TGTCCTTTCC 1560  
 10 TAATAAAATG AGGAAATTGC ATCGCATTGT CTGAGTAGGT GTCATTCTAT TCTGGGGGGT 1620  
 GGGGTGGGGC AGGACAGCAA GGGGGAGGAT TGGGAAGACA ATAGCAGGCA TGCTGGGGAT 1680  
 GCGGTGGGCT CTATGGAACC AGCTGGGGCT CGACAGCTAT GCCAAGTACG CCCCCATTG 1740  
 15 ACGTCAATGA CGGTAAATGG CCCGCTGGC ATTATGCCA GTACATGACC TTATGGGACT 1800  
 TTCCTACTTG GCAGTACATC TACGTATTAG TCATCGCTAT TACCATGGTG ATCGGTTTTT 1860  
 GGCAGTACAT CAATGGGCGT GGATAGCGT TTAGCTCACG GGGATTTCCA AGTCTCCACC 1920  
 CCATTGACGT CAATGGGAGT TTGTTTTGGC ACCAAAATCA ACGGGACTTT CCAAAATGTC 1980  
 20 GTAACAATC CGCCCCATTG ACGCAAATGG GCGGTAGGCG TGTACGGTGG GAGGTCTATA 2040  
 TAAGCAGAGC TGGGTACGTC CTCACATTCA GTGATCAGCA CTGAACACAG ACCCGTCGAC 2100  
 ATGGGTTGGA GCCTCATCTT GCTCTTCCTT GTCGCTGTG CTACGCGTGT CGCTAGCACC 2160  
 25 AAGGGCCCAT CGGTCTTCCC CTTGGCACCC TCCTCCAAGA GCACCTCTGG GGGCACAGCG 2220  
 GCCCTGGGCT GCCTGGTCAA GGACTIONC CCCGAACCGG TGACGGTGTG GTGGAATCA 2280  
 GCGCCCTGA CCAGCGGCGT GCACACCTTC CCGGCTGTCC TACAGTCCTC AGGACTCTAC 2340  
 30 TCCCTCAGCA GCGTGGTGAC CGTGCCCTCC AGCAGCTTGG GCACCCAGAC CTACATCTGC 2400  
 AACGTGAATC ACAAGCCCAG CAACACCAAG GTGGACAAGA AAGCAGAGCC CAAATCTTGT 2460  
 GACAAAATC ACACATGCCC ACCGTGCCCC GCACCTGAAC TCCTGGGGGG ACCGTCAGTC 2520  
 35 TTCTCTTCC CCCCCAAACC CAAGGACACC CTCATGATCT CCCGGACCCC TGAGGTCA 2580  
 TGCGTGGTGG TGGACGTGAG CCACGAAGAC CTTGAGGTCA AGTTCAACTG GTACGTGGAC 2640  
 GCGTGGAGG TGCATAATGC CAAGACAAAG CCGGGGAGG AGCAGTACAA CAGCACGTAC 2700  
 40 CGTGTGGTCA GCGTCCTCAC CGTCCTGCAC CAGGACTGGC TGAATGGCAA GGAGTACAAG 2760  
 TGCAAGGTCT CCAACAAAGC CCTCCAGCC CCCATCGAGA AAACCATCTC CAAAGCCAAA 2820  
 GGGCAGCCCC GAGAACCACA GGTGTACACC CTGCCCCCA? CCCGGGATGA GCTGACCAAG 2880  
 45 AACCAGGTCA GCCTGACCTG CTTGGTCAA GCTTCTATC CCAGCGACAT CGCCGTGGAG 2940  
 TGGGAGAGCA ATGGGCAGCC GGAGAACAAC TACAAGACCA CGCCTCCCGT GCTGGACTCC 3000  
 GACGGCTCCT TCTTCTCTA CAGCAAGCTC ACCGTGGACA AGAGCAGGTG GCAGCAGGGG 3060  
 50 AACGTCTTCT CATGCTCCGT GATGCATGAG GCTCTGCACA ACCACTACAC GCAGAAGAGC 3120  
 CTCTCCCTGT CTCCGGGTAA ATGAGGATCC GTTAACGGTT ACCAACTACC TAGACTGGAT 3180

55

TCGTGACAAC ATGCGGCCGT GATATCTACG TATGATCAGC CTCGACTGTG CCTTCTAGTT 3240  
 GCCAGCCATC TTTTGTGTTG CCTTCCCCCG TGCCTTCCTT GACCCTGGAA GGTGCCACTC 3300  
 5 CCACTGTCTT TTTCTAATAA AATGAGGAAA TTGCATCGCA TTGTCTGAGT AGGTGTCATT 3360  
 CTATCTCTGG GGTGGGGTG GGGCAGGACA GCAAGGGGGA GGATTGGGAA GACAATACCA 3420  
 GGCATGCTGG GGATGCGGTG GGCTCTATGG AACCAGCTGG GGCTCGACAG CGCTGGATCT 3480  
 10 CCCGATCCCC AGCTTTGCTT CTCAATTTCT TATTTGCATA ATGAGAAAAA AAGGAAAATT 3540  
 AATTTTAACA CCAATTCAGT AGTTGATTGA GCAAATGCGT TGCCAAAAAG GATGCTTTAG 3600  
 AGACAGTGTT CTCTGCACAG ATAAGGACAA ACATTATTCA GAGGGAGTAC CCAGAGCTGA 3660  
 15 GACTCCTAAG CCAGTGAGTG GCACAGCATT CTAGGGAGAA ATATGCTTGT CATCACCAGAA 3720  
 GCCTGATTCC GTAGAGCCAC ACCTTGGTAA GGGCCAATCT GCTCACACAG GATAGAGAGG 3780  
 GCAGGAGCCA GGCAGAGCA TATAAGGTGA GGTAGGATCA GTTGCTCCTC ACATTTGCTT 3840  
 20 CTGACATAGT TGTGTTGGGA GCTTGGATAG CTTGGACAGC TCAGGGCTGC GATTTCCGCG 3900  
 CAAACTTGAC GGCAATCCTA GCGTGAAGGC TGGTAGGATT TTATCCCCGC TGCCATCATG 3960  
 GTTCGACCAAT TGAAGTGCAT CGTCGCCGTG TCCCAAAATA TGGGGATTGG CAAGAACGGA 4020  
 25 GACCTACCCT GGCCTCCGCT CAGGAACGAG TTCAAGTACT TCCAAAGAAT GACCACAACC 4080  
 TCTTCAGTGG AAGGTAAACA GAATCTGGTG ATTATGGGTA GGAAAACCTG GTTCTCCATT 4140  
 CCTGAGAAGA ATCGACCTTT AAAGGACAGA ATTAATATAG TTCTCAGTAG AGAACTCAA 4200  
 30 GAACCACCAC GAGGAGCTCA TTTTCTTGCC AAAAGTTTGG ATGATGCCTT AAGACTTATT 4260  
 GAACAACCGG AATTGGCAAG TAAAGTAGAC ATGGTTTGGG TAGTGGGAGG CAGTTCTGTT 4320  
 TACCAGGAAG CCATGAATCA ACCAGGCCAC CTTAGACTCT TTGTGACAAG GATCATGCAG 4380  
 35 GAATTTGAAA GTGACACGTT TTTCCAGAA ATTGATTTGG GGAAATATAA ACTTCTCCCA 4440  
 GAATACCCAG GCGTCCTCTC TGAGGTCCAG GAGGAAAAAG GCATCAAGTA TAAGTTTGAA 4500  
 GTCTACGAGA AGAAAGACTA ACAGGAAGAT GCTTTCAAGT TCTCTGCTCC CCTCCTAAAG 4560  
 40 CTATGCATTT TTATAAGACC ATGGGACTTT TGCTGGCTTT AGATCAGCCT CGACTGTGCC 4620  
 TTCTAGTTGC CAGCCATCTG TTGTTTGCCC CTCCCCGTG CCTTCCTTGA CCCTGGAAGG 4680  
 TGCCACTCCC ACTGTCCTTT CTAATAAAA TGAGGAAATT GCATCGCATT GTCTGAGTAG 4740  
 45 GTGTCATTCT ATTCTGGGGG GTGGGGTGGG GCAGGACAGC AAGGGGGAGG ATTGGGAAGA 4800  
 CAATAGCAGG CATGCTGGGG ATGCGGTGGG CTCTATGGAA CCAGCTGGGG CTCGAGCTAC 4860  
 TAGCTTTGCT TCTCAATTC TTATTTGCAT AATGAGAAAA AAAGGAAAAT TAATTTTAAC 4920  
 50 ACCAATTCAG TAGTTGATTG AGCAAATGCG TTGCCAAAA GGATGCTTTA GAGACAGTGT 4980  
 TCTCTGCACA GATAAGGACA AACATTATTC AGAGGGAGTA CCCAGAGCTG AGACTCCTAA 5040

55

GCCAGTGAGT GGCACAGCAT TCTAGGGAGA AATATGCTTG TCATCACCAG AGCCTGATTC 5100  
 CGTAGAGCCA CACCTTGGTA AGGGCCAATC TGCTCACACA GGATAGAGAG GGCAGGAGCC 5160  
 5 AGGGCAGAGC ATATAAGGTG AGGTAGGATC AGTTGCTCCT CACATTTGCT TCTGACATAG 5220  
 TTGTGTTGGG AGCTTGGATC GATCCTCTAT GGTGAACAA GATGGATTGC ACGCAGGTTC 5280  
 TCCGGCCGCT TGGCTGGAGA GGCTATTCGG CTATGACTGG GCACAACAGA CAATCGGCTG 5340  
 10 CTCTGATGCC GCGTGTTC GGTGTTCAGC GCAGGGGCGC CCGGTTCTTT TTGTCAAGAC 5400  
 CGACCTGTCC GGTGCCCTGA ATGAAGTGA GGACGAGGCA GCGCGGCTAT CGTGGCTGGC 5460  
 CACGACGGGC GTTCCTTGGC CAGCTGTGCT CAGCTTTGTC ACTGAAGCGG GAAGGGACTG 5520  
 15 GCTGCTATTG GCGAAGTGC CGGGGCAAGA TCTCTGTGA TCTACCTTG CTCCTGCCGA 5580  
 GAAAGTATCC ATCATGGCTG ATGCAATGCG GCGGCTGCAT ACGCTTGATC CGGCTACCTG 5640  
 CCCATTGAC CACCAAGCGA AACATCGCAT CAGGCGAGCA CGTACTCGGA TGGAAAGCCG 5700  
 20 TCTTGTGAT CAGGATGATC TGGACGAAGA GCATCAGGGG CTCGCGCCAG CCGAAGTGT 5760  
 CGCCAGGCTC AAGGCGCGCA TGGCCGACGG CAGGATCTC CTCGTGACCC ATGGCGATGC 5820  
 CTGCTTGCCG AATATCATGG TGGAAATGG CCGCTTTCTT GGATTCATCG ACTGTGGCCG 5880  
 25 GCTGGGTGTG CCGGACCGCT ATCAGGACAT AGCGTTGGCT ACCCGTGATA TTGCTGAAGA 5940  
 GCTTGGCGCC GAATGGGCTG ACCGCTTCCT CGTGCTTAC GGTATCGCCG CTCCTGATTC 6000  
 GCAGCGCATC GCCTTCTATC GCCTTCTGA GAGTCTTC TGAGCGGAC TGTGGGTTC 6060  
 30 GAAATGACCG ACCAAGCGAC GCCCAACCTG CCATCAGAG ATTTGATTC CACCGCCGCC 6120  
 TTCTATGAA GGTGGGCTT CGGAATGCTT TTCCGGGACG CCGCTGGAT GATCCTCCAG 6180  
 CGCGGGGATC TCATGCTGGA GTTCTTCGCC CACCCCACT TGTTTATTGC AGCTTATAAT 6240  
 35 GGTACAAAT AAAGCAATAG CATCACAAT TTCACAATA AAGCATTTT TTCACTGCAT 6300  
 TCTAGTTGTG GTTTGTCCA ACTCATCAAT CTATCTTATC ATGTCTGGAT CGCGGCCGCG 6360  
 ATCCCGTCGA GAGCTTGGCG TAATCATGGT CATAGCTGTT TCCTGTGTGA AATTGTTATC 6420  
 CGCTCACAAT TCCACACAAC ATACGAGCCG GAAGCATAAA GTGTAAAGCC TGGGGTGCCCT 6480  
 40 AATGAGTGAG CTAATCACA TTAATTGCGT TGCGCTCACT GCCCGCTTTC CAGTCGGGAA 6540  
 ACCTGTCGTG CCAGCTGCAT TAATGAATCG GCCAACCGC GGGGAGAGGC GGTTTGCGTA 6600  
 TTGGGCGCTC TTCGCTTCC TCCTCACTG ACTCGCTGCG CTCGCTCGTT CGGCTGCGGC 6660  
 45 GAGCCGTATC AGCTCACTCA AAGCGGTAA TACGGTTATC CACAGAATCA GGGGATAACG 6720  
 CAGGAAAGAA CATGTGAGCA AAAGCCGAG AAAGCCGAG GAACCGTAAA AAGGCCGCT 6780  
 TGCTGGCGTT TTCCATAGG CTCGCGCCCG CTGACGAGCA TCACAAAAAT CGACGCTCA 6840  
 50 CTCAGAGGTG GCGAAACCG ACAGGACTAT AAAGATACCA GCGCTTTCCC CCTGGAAGCT 6900  
 CCCTCGTGG CTCTCTGTT CCGACCTGC CGCTTACCG ATACCTGTCC GCCTTCTCC 6960

55

	CTTCGGGAAC CGTGGCGCTT TCTCAATGCT CACGCTGTAG GTATCTCAGT TCGGTGTAGG	7020
	TCGTTCCGTC CAAGCTGGGC TGTGTGCACG AACCCCGCGT TCAGCCCGAC CGCTGGCGCT	7080
5	TATCCGGTAA CTATCGTCTT GAGTCCAACC CGGTAAGACA CGACTTATCG CCACTGGCAG	7140
	CAGCCACTGG TAACAGGATT AGCAGAGCGA GGTATGTAGG CGGTGCTACA GAGTTCTTGA	7200
	AGTGGTGGCC TAACTACGGC TACACTAGAA GCACAGTATT TGGTATCTGC GCTCTGCTGA	7260
10	AGCCAGTTAC CTTCGGAAAA AGAGTTGGTA GCTCTGATC CGGCAAAACA ACCACCGCTG	7320
	GTAGCGGTGG TTTTTTGTG TCAAGCAGC AGATTACGGC CAGAAAAAA GGATCTCAAG	7380
	AAGATCCTTT GATCTTTTCT ACGGGGTCTG ACGCTCAGTG GAACGAAAAC TCACGTAAAG	7440
15	GGATTTTGGT CATGAGATTA TCAAAAAGGA TCTTCACCTA GATCCTTTTA AATTAAAAAT	7500
	GAAGTTTAA ATCAATCTAA AGTATATATG AGTAAACTTG GTCTGACAGT TACCAATGCT	7560
	TAATCACTGA GGCACCTATC TCAGCGATCT GTCTATTTCT TTCATCCATA GTTGCCTGAC	7620
20	TCCTCCGTCT GTAGATAACT ACQATACGGG AGGGCTTACC ATCTGGCCCC AGTGCTGCAA	7680
	TGATACCCGG AGACCCACGC TCACCCGGCTC CAGATTTATC AGCAATAAAC CAGCCAGCCG	7740
	GAAGGSCCGA GCGCAGAAGT GGTCTCTCAA CTTTATCCGC CTCCATCCAG TCTATTAATT	7800
25	GTTGCCGGGA AGCTAGAGTA AGTAGTTCCG CAGTTAATAG TTTGCCCAAC GTTGTGCCA	7860
	TTGCTACAGG CATCGTGGTG TCACGCTCGT CGTTTGGTAT GGCTTCATTC AGCTCCGGTT	7920
	CCCAACGATC AAGGCGAGTT ACATGATCCC CCATGTTGTG CAAAAAAGCG GTTAGCTCCT	7980
30	TCGGTCCCTCC GATCGTTGTC AQAAGTAAGT TGCCCGCAGT GTTATCACTC ATGTTTATCG	8040
	CAGCACTGCA TAATTCTCTT ACTGTCATGC CATCCGTAAG ATGCTTTTCT GTGACTGGTG	8100
	AGTACTCAAC CAAGTCATTC TGAQAATAGT GTATCGGCG ACCGAGTTGC TCTTGCCCGG	8160
35	CGTCAATACG GDATAATACC GCQCCACATA GCQAACTTT AAAAGTGCTC ATCATTGGAA	8220
	AACGTTCTTC CCGGCGAAAA CTCTCAAGGA TCTTACCGCT GTTGAGATCC AGTTGATGT	8280
	AACCCACTCG TGCACCCAAC TGATCTTCAG CATCTTTTAC TTTCACCAGC GTTCTGGGT	8340
40	GAGCAAAAAC AGGAAGGCAA AATGCCGCAA AAAAGGGAAT AAGGGCGACA CGGAAATGTT	8400
	GAATACTCAT ACTCTTCCTT TTTCAATATT ATTGAAGCAT TTATCAGGGT TATTGTCTCA	8460
	TGAGCGGATA CATATTTGAA TGTATTTAGA AAAATAAACA AATAGGGGT CCGCGCACAT	8520
45	TTCCCCGAAA AGTGCCACCT	8540

## (3) INFORMATION FOR SEQ ID NO: 2:

## (i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 9209 bases

(B) TYPE: nucleic acid  
 (C) STRANDEDNESS: single  
 (D) TOPOLOGY: circular

(ii) MOLECULE TYPE: DNA (genomic)

(iii) HYPOTHETICAL: yes

(iv) ANTI-SENSE: no

(ix) SEQUENCE DESCRIPTION: SEQ ID NO: 2:

15	GACGTCGCGG CCGCTCTAGG CCTCCAAAAA AGCCTCCTCA CTACTTCTGG AATAGCTCAG	60
	AGGCCGAGGC GGCCTCGGCC TCTGCATAAA TAAAAAAAT TAGTCAGCCA TGCATGGGGC	120
	GGAGAATGGG CGGAACTGGG CGGAGTTAGG GGCGGGATGG GCGGAGTTAG GGGCGGGACT	180
20	ATGGTTGCTG ACTAATTGAG ATGCATGCTT TGCATACTTC TGCCTGCTGG GGAGCCTGGG	240
	GACTTTCCAC ACCTGGTTGC TGAATAATTG AGATGCATGC TTTGCATACT TCTGCCTGCT	300
	GGGGAGCCTG GGGACTTTCC ACACCCTAAC TGACACACAT TCCACAGAAT TAATTCCCCT	360
25	AGTTATTAAT AGTAATCAAT TACGGGGTCA TTAGTTCATA GCCCATATAT GGAGTTCCGC	420
	GTTACATAAC TTACGGTAAA TGGCCCGCCT GGCTGACCGC CCAACGACCC CCGCCCATTG	480
	ACCTCAATAA TGACGTATGT TCCCATAGTA ACGCCAATAG GGACTTTCCA TTGACGTCAA	540
30	TGGGTGGACT ATTTACGGTA AACTGCCCCAC TTGGCAGTAC ATCAAGTGTA TCATATGCCA	600
	AGTACGCCCC CTATTGACGT CAATGACGGT AAATGGCCCG CCTGGCATTG TGGCCAGTAC	660
	ATGACCTTAT GGGACTTTCC TACTTGCCAG TACTCTACG TATTAGTCAT CGCTATTACC	720
	ATGGTGATGC GGTTTTGGCA GTACATCAAT GGGCGTGGAT AGCGGTTTGA CTCACGGGGA	780
35	TTTCCAAGTC TCCACCCCAT TGACGTCAAT GGGAGTTTGT TTTGGCACCA AAATCAACGG	840
	GACTTTCCAA AATGTCGTAA CAACTCCGCC CCATTGACGC AAATGGGCGG TAGGCGTGTA	900
	CGGTGGGAGG TCTATATAAG CAGAGCTGGG TACGTGAACC GTCAGATCGC CTGGAGACGC	960
40	CATCACAGAT CTCTCACTAT GGATTTTCAG GTGCAGATTA TCAGCTTCCT GCTAATCAGT	1020
	GCTTCAGTCA TAATGTCCAG AGGACAAATT GTTCTCTCCC AGTCTCCAGC AATCCTGTCT	1080
	GCATCTCCAG GGGAGAAGGT CACAATGACT TGCAGGGCCA GCTCAAGTGT AAGTTACATC	1140
45	CACTGGTTCC AGCAGAAGCC AGGATCCTCC CCCAAACCCT GGATTATATGC CACATCCAAC	1200
	CTGGCTTCTG GAGTCCCTGT TCGCTTCAGT GGCAGTGGGT CTGGGACTTC TTA CTCTCTC	1260
	ACAATCAGCA GAGTGGAGGC TGAAGATGCT GCCACTTATT ACTGCCAGCA GTGGACTAGT	1320
50	AACCCACCCA CGTTCGGAGG GGGGACCAAG CTGGAAATCA AACGTACGGT GGCTGCACCA	1380
	TCTGTCTTCA TCTTCCCGCC ATCTGATGAG CAGTTGAAAT CTGGAAGTGC CTCTGTTGTG	1440



TGCCTGCTGA ATAACTTCTA TCCCAGAGAG GCCAAAGTAC AGTGGAAGGT GGATAACGCC 1500  
 CTCCAATCGG GTAACCTCCA GGAGAGTGTC ACAGAGCAGG ACAGCAAGGA CAGCACCTAC 1560  
 5 AGCCTCAGCA GCACCCTGAC GCTGAGCAAA GCAGACTACG AGAAACACAA AGTCTACGCC 1620  
 TGCGAAGTCA CCCATCAGGG CCTGAGCTCG CCCGTCACAA AGAGCTTCAA CAGGGGAGAG 1680  
 TGTTGAAATC AGATCCGTTA ACGGTTACCA ACTACCTAGA CTGGATTCTG GACAACATCC 1740  
 10 GGCCGTGATA TCTACGTATG ATCAGCCTCG ACTGTGCCTT CTAGTTGCCA GCCATCTGTT 1800  
 GTTTGCCCCCT CCCCCGTGCC TTCCTTGACC CTGGAAGGTG CCACTCCCAC TGTCTTTCC 1860  
 TAATAAAATG AGGAAATTGC ATCGCATTGT CTGAGTAGGT GTCATTCTAT TCTGGGGGGT 1920  
 15 GGGGTGGGGC AGGACAGCAA GGGGGAGGAT TGGGAAGACA ATAGCAGGCA TGCTGGGGAT 1980  
 GCGGTGGGCT CTATGGAACC AGCTGGGGCT CGACAGCTAT GCCAAGTACG CCCCCTATTG 2040  
 ACGTCAATGA CGGTAAATGG CCCGCCTGGC ATTATGCCCA GTACATGACC TTATGGGACT 2100  
 20 TTCCTACTTG GCAGTACATC TACGTATTAG TCATCGCTAT TACCATGGTG ATGCGGTTTT 2160  
 GGCAGTACAT CAATGGGCGT GGATAGCGGT TTGACTCAGC GGGATTTCCTA AGTCTCCACC 2220  
 CCATTGACGT CAATGGGAGT TTGTTTGGC ACCAAAATCA ACGGGACTTT CCAAAATGTC 2280  
 25 GTAACAACTC CGCCCCATTG ACGCAAATGG GCGGTAGGCG TGTACGGTGG GAGGTCTATA 2340  
 TAAGCAGAGC TGGGTACGTC CTCACATTCA GTGATCAGCA CTGAACACAG ACCCGTCGAC 2400  
 ATGGGTGGA GCCTCATCTT GCTCTTCCTT GTCGCTGTG CTACGCGTGT CCTGTCCCAG 2460  
 30 GTACAACTGC AGCAGCCTGG GGCTGAGCTG GTGAAGCCTG GGGCCTCAGT GAAGATGTCC 2520  
 TGCAAGGCTT CTGGCTACAC ATTTACCAGT TACAATATGC ACTGGGTAAA ACAGACACCT 2580  
 GGTGCGGGCC TGAATGGAT TGGAGCTATT TATCCCGGAA ATGGTGATAC TTCCTACAAT 2640  
 35 CAGAAGTTCA AAGGCAAGGC CACATTGACT GCAGACAAAT CCTCCAGCAC AGCCTACATG 2700  
 CAGCTCAGCA GCCTGACATC TGAGGACTCT GCGGTCTATT ACTGTGCAAG ATCGACTTAC 2760  
 TACGGCGGTG ACTGGTACTT CAATGTCTGG GGCGCAGGGA CCACGGTCAC CGTCTCTGCA 2820  
 40 GCTAGCACCA AGGGCCCATC GGTCTTCCCC CTGGCACCCCT CCTCCAAGAG CACCTCTGGG 2880  
 GGCACAGCGG CCCTGGGCTG CCTGGTCAAG GACTACTTCC CCGAACCAGT GACGGTGTGG 2940  
 TGGAACCTCAG GCGCCCTGAC CAGCGGCGTG CACACCTTCC CGGCTGTCTT ACAGTCTTCA 3000  
 GGACTCTACT CCCTCAGCAG CGTGGTGACC GTGCCCTCCA GCAGCTTGGG CACCCAGACC 3060  
 45 TACATCTGCA ACGTGAATCA CAAGCCCAGC AACACCAAGG TGGACAAGAA AGCAGAGCCC 3120  
 AAATCTTGTG ACAAACCTCA CACATGCCCA CCGTGCCAG CACCTGAACT CCTGGGGGGA 3180  
 CCGTCAGTCT TCCTCTTCCC CCAAAAACCC AAGGACACCC TCATGATCTC CCGGACCCCT 3240  
 50 GAGGTCACAT GCGTGGTGGT GGACGTGAGC CACGAAGACC CTGAGGTCAA GTTCAACTGG 3300  
 TACGTGGACG GCGTGGAGGT GCATAATGCC AAGACAAAGC CGCGGGAGGA GCAGTACAAC 3360

55

AGCACGTACC GTGTGGTCAG CGTCCTCACC GTCCTGCACC AGGACTGGCT GAATGGCAAAG 3420  
 GAGTACAAGT GCAAGGTCTC CAACAAAGCC CTCCCAGCCC CCATCGAGAA AACCATCTCC 3480  
 5 AAAGCCAAAG GGCAGCCCCG AGAACCACAG GTGTACACCC TGCCCCCCTC CCGGGATGAG 3540  
 CTGACCAACA ACCAGGTCAG CCTGACCTGC CTGGTCAAAG GCTTCTATCC CAGCGACATC 3600  
 GCGGTGGAGT GCGAGAGCAA TGGGCAGCCG GAGAACAAC ACAAAGACCAC GCCTCCCGTG 3660  
 10 CTGGACTCCG ACSGCTCCTT CTTCCTCTAC AGCAAGCTCA CCGTGGACAA GAGCAGGTGG 3720  
 CAGCAGCGGA ACCTCTTCTC ATGCTCCGTG ATGCATGAGG CTCTGCACAA CCACTACACG 3780  
 CAGAAGAGCC TCTCCCTGTC TCCGGCTAAA TGAGGATCCG TTAACGGTTA CCAACTACCT 3840  
 15 AGACTGGATT CSTGACAACA TGCGGCCCGT ATATCTACGT ATGATCAGCC TCGACTGTGC 3900  
 CTCTAGTTG CCGCCATCT GTTGTTCGCC CCTCCCCCGT GCCTTCCTTG ACCCTGGAAG 3960  
 GTGCCACTCC CACTCTCCTT TCCTAATAAA ATGAGGAAAT TGATCCCAT TGTCTGAGTA 4020  
 20 GGTGTCATTC TATCTGGGG GGTGGGGTGG GGCAGGACAG CAAGGGGGAG GATTGGGAAG 4080  
 ACAATACCAG GCATGCTGGG GATGCGGTGG GCTCTATGGA ACCAGCTGGG GCTCGACAGC 4140  
 GCTGGATCTC CCGATCCCCA GCTTTGCTTC TCAATTCTT ATTTGCATAA TGAGAAAAAA 4200  
 25 AGGAAAATTA ATTTAAACAC CAATTCAGTA GTTGATTGAG CAAATCCGTT GCCAAAAAGG 4260  
 ATGCTTTAGA GACAGTGTTC TCTGCACAGA TAAGGACAAA CATTATTCAG AGGGAGTACC 4320  
 CAGAGCTGAG ACTCCTAAGC CAGTGAGTGG CACAGCATTC TAGGGAGAAA TATGCTTGTC 4380  
 30 ATCACCGAAG CCTGATTCCG TAGAGCCACA CCTTGGTAAG GGCCAATCTG CTCACACAGG 4440  
 ATAGACAGGG CAGGAGCCAG GGCAGAGCAT ATAAGGTGAG GTAGGATCAG TTGCTCCTCA 4500  
 CATTTCCTTC TGACATAGTT GTCTTGGGAG CTTCGATAGC TTGGACAGCT CAGGGCTGCC 4560  
 35 ATTTCCGCCC AAACCTGACG GCAATCCTAG COTGAAGCCT GGTAGGATTT TATCCCCGCT 4620  
 GCCATCATCG TTCGACCATT GAACTGCATC GTCGCCGTCT CCCAAAATAT GGGGATTGGC 4680  
 AAGAACGGAG ACCTACCCTG GCCTCCGCTC AGGAACGAGT TCAAGTACTT CCAAAGAATG 4740  
 40 ACCACAACCT CTTCAAGTGA AGCTAAACAG AATCTGGTGA TTATGGGTAG GAAAACCTGG 4800  
 TTCTCCATTC CTGAGAAGAA TCGACCTTGA AAGGACAGAA TTAATATAGT TCTCAGTAGA 4860  
 GAACTCAAAG AACCACCACG AGGAGCTCAT TTTCTTGCCA AAAGTTTGGG TGATGCCTTA 4920  
 45 AGACTTATTG AACAAACCGA ATTGGCAAGT AAAGTAGACA TCGTTTGGAT AGTCGGAGGC 4980  
 AGTTCTGTTT ACCAGGAAGC CATGAATCAA CCAGGCCACC TTAGACTCTT TGTGACAAGG 5040  
 ATCATGCAGG AATTTGAAAG TGACACGTTT TTCCAGAAA TTGATTTGGG GAAATATAAA 5100  
 50 CTTCTCCCAG AATACCCAGG CGTCTCTCT GAGCTCCAGG ACCAAAAAGG CATCAAGTAT 5160  
 AAGTTTGAAG TCTACGAGAA GAAAGACTAA CAGGAAGATG CTTTCAAGTT CTCTGCTCCC 5220

CTCTAAAGC TATGCATTTT TATAAGACCA TGGGACTTTT GCTGGCTTTA GATCAGCCTC 5280  
 GACTGTGCCT TCTAGTTGCC AGCCATCTGT TGTGTGCCCC TCCCCCGTGC CTTCCCTTGAC 5340  
 5 CCTGGAAGGT GCCACTCCCA CTGTCCTTTC CTAATAAAAT GAGGAAATTG CATCGCATTG 5400  
 TCTGAGTAGG TGTCATTCTA TTCTGGGGGG TGGGGTGGGG CAGGACAGCA AGGGGGAGGA 5460  
 TTGGGAAGAC AATAGCAAGC ATGCTGGGGA TGGGUTGGGC TCTATGGAAC CAUGTGGGGC 5520  
 10 TCGAGCTACT AGCTTTGCTT CTCAATTCTT TATTTGCATA ATGAGAAAAA AAGGAAAATT 5580  
 AATTTTAACA CCAATTCAGT AGTTGATTGA GCAAAATGCGT TGCCAAAAAG GATGCTTTAG 5640  
 AGACAGTGTT CTCTGCACAG ATAAGGACAA ACATTATTCA GAGGGAGTAC CCAGAGCTGA 5700  
 15 GACTCCTAAG CCAGTGAGTG GCACAGCATT CTAGGGAGAA ATATGCTTGT CATCACCGAA 5760  
 GCCTGATTCC GTAGAGCCAC ACCTTGGTAA GGGCCAATCT GCTCACACAG GATAGAGAGG 5820  
 GCAGGAGCCA GGGCAGAGCA TATAAGGTGA GGTAGGATCA GTTGCTCCTC ACATTTGCTT 5880  
 20 CTGACATAGT TGTGTTGGGA GCTTGGATCG ATCCTCTATG GTTGAACAAG ATGGATTGCA 5940  
 CGCAGGTTCT CCGGCCGCTT GGGTGGAGAG GCTATTGCGC TATGACTGGG CACAACAGAC 6000  
 AATCGGCTGC TCTGATGCCG CCGTGTTCGG GCTGTGAGCG CAGGGGCGCC CGGTTCTTTT 6060  
 25 TGTCAAGACC GACCTGTCCG GTGCCCTGAA TGAAGTGCAG GACGAGGCAG CGCGGCTATC 6120  
 GTGGCTGGCC ACGACGGGCG TTCCTTGCGC AGCTGTGCTC GACGTTGTCA CTGAAGCGGG 6180  
 AAGGGACTGG CTGCTATTGG GCGAAGTGCC GGGGCAGGAT CTCCTGTCAT CTCACCTTGC 6240  
 30 TCCTGCCGAG AAAGTATCCA TCATGGCTGA TGCAATGCGG CGGCTGCATA CGCTTGATCC 6300  
 GGCTACCTGC CCATTGACCC ACCAAGCGAA ACATCGCATC GAGCGAGCAC GTACTCGGAT 6360  
 GGAAGCCGGT CTTGTGATC AGGATGATCT GGACGAAGAG CATCAGGGGC TCGCGCCAGC 6420  
 CGAACTGTTT GCCAGGCTCA AGGCGCGCAT GCGCGACGGC GAGGATCTCG TCGTGACCCA 6480  
 35 TGGCGATGCC TGCTTGCCGA ATATCATGGT GGAAATGGC CGCTTTTCTG GATTCATCGA 6540  
 CTGTGGCCGG CTGGGTGTGG CGGACCGCTA TCAGGACATA GCGTTGGCTA CCCGTGATAT 6600  
 TGCTGAAGAG CTTGGCGGCG AATGGGCTGA CCGCTTCCTC GTGCTTTACG GTATCGCCGC 6660  
 40 TCCCGATTCT CAGCGCATCG CCTTCTATCG CCTTCTTGAC GAGTTCTTCT GAGCGGGACT 6720  
 CTGGGGTTCT AAATGACCGA CCAAGCGACG CCCAACCTGC CATCACGAGA TTTCGATTCC 6780  
 ACCGCCGCCT TCTATGAAAG GTTGGGCTTC GGAATCGTTT TCCGGGACGC CGGCTGGATG 6840  
 45 ATCTCCAGC GCGGGGATCT CATGCTGGAG TTCTTCGCCC ACCCCAATT GTTTATTGCA 6900  
 GCTTATAATG GTTACAAATA AAGCAATAGC ATCACAAATT TCACAAATAA AGCATTTTTT 6960  
 TCACTGCATT CTAGTTGTGG TTTGTCCAAA CTCATCAATC TATCTTATCA TGTCTGGATC 7020  
 50 GCGGCCGCGA TCCCGTCGAG AGCTTGGCGT AATCATGGTC ATAGCTGTTT CCTGTGTGAA 7080  
 ATTGTTATCC GCTCACAATT CCACACAACA TACGAGCCGG AAGCATAAAG TGTAAGCCT 7140

55

5  
 10  
 15  
 20  
 25  
 30  
 35  
 40  
 45  
 50  
 55

GGGGTGCGTA ATGAGTGAGC TAACTCACAT TAATTGCGTT GCGCTCACTG CCCGTTTTCC 7200  
 AGTCGGGAAA CCTGTCGTGC CAGCTGCATT AATGAATCGG CCAACGGCGG GGGAGAGGCG 7260  
 GTTTGCGTAT TGGCGCGTCT TCCGCTTCCT CGCTCACTGA CTCGCTGCGC TCGGTGCTTC 7320  
 GCGTGCGGCG AGCGGTATCA GCTCACTCAA AGCGGTAAT ACGGTTATCC ACAGAATCAG 7380  
 GGGATACGCG AGGAAAGAAC ATGTGAGCAA AAGGCCAGCA AAAGGCCAGG AACCGTAAAA 7440  
 AGCGCGCGTT GCTGGCGTTT TTCCATAGGC TCGCGCCCCC TCACGAGCAT CACAAAAATC 7500  
 GACGCTCAAG TCAGAGGTGG CGAAACCCGA CAGGACTATA AAGATACCAQ GCGTTTCCCC 7560  
 CTGGAAGCTC CCTCGTGCGC TCTCCTGTTT CGACCCCTGCC GCTTACCGGA TACCTGTCCG 7620  
 CTTTCTCCCC TTCGGGAAGC GTGGCGCTTT CTCAATGCTC ACGCTGTAGG TATCTCAGTT 7680  
 CGGTGTAGCT CTTTCGCTCC AAGCTGGGCT GTGTGCACGA ACCCCCCGTT CAGCCCGACC 7740  
 GCTGCGCCTT ATCCGGTAAC TATCGTCTTG AGTCCAACCC GGTAAAGACAC GACTTATCGC 7800  
 CACTGGCAGC AGCCACTGGT AACAGGATTA CCAGAGCGAC GTATGTAGGC GGTGCTACAG 7860  
 AGTTCTTGAA GTGGTGGCCT AACTACGGCT AACTAGAAG GACAGTATTT GGTATCTGCG 7920  
 CTCTGCTGAA GCCAGTTACC TTCGGAAAAA GAGTTGCTAG CTCTTGATCC GCGAAACAAA 7980  
 CCACCGCTGG TAGCGGTGGT TTTTGTGTTT GCAAGCAGCA GATTACGCGC AGAAAAAAG 8040  
 GATCTCAAGA AGATCCTTTG ATCTTTTCTA GGGGCTCTGA CGCTCAGTGG AACGAAAACT 8100  
 CACGTTAAGG GATTTTGGTC ATGAGATTAT CAAAAAGGAT CTTACCTAG ATCCTTTTAA 8160  
 ATTAAAAATG AAGTTTTTAA TCAATCTAAA GTATATATGA GTAAACTTGG TCTGACAGTT 8220  
 ACCAATGCTT AATCAGTGAG GCACCTATCT CAGCGATCTG TCTATTTCTG TCATCCATAG 8280  
 TTGCCTGACT CCCCCTCGTG TAGATAACTA CGATACGGGA GGGCTTACCA TCTGGCCCCA 8340  
 GTGCTGCAAT GATACCGCGA GACCCACGCT CACCGGCTCC AGATTATCA GCAATAAACG 8400  
 AGCCAGCCGG AAGGGCCGAG CGCAGAACTG GTCCCTGCAC TTTATCCGCG TCCATCCAGT 8460  
 CTATTAATTG TTGCCGGGAA GCTAGAGTAA GTAGTTCCGC AGTTAATACT TTGCCAACC 8520  
 TTGTTGCCAT TGCTACAGGC ATCGTGGTGT CACGCTCGTC GTTTGGTATG GCTTCATTCA 8580  
 GCTCCGGTTC CCAACGATCA AGGCGAGTTA CATGATCCCC CATGTTCTGC AAAAAAGCGG 8640  
 TTAGCTCCTT CGGTCTCTCG ATCGTTGTCA GAAGTAACTT GGCCGCAGTG TTATCACTCA 8700  
 TGGTTATGGC AGCACTGCAT AATTCTCTTA CTGTCAAGCC ATCCGTAAGA TGCTTTTCTG 8760  
 TGA CTGGTGA GTACTCAACC AAGTCATTCT GAGAATAGTG TATGCGGCGA CCGAGTTGCT 8820  
 CTTGCCCCGC GTCAATACGG GATAATACCG CGCCACATAG CAGAACTTTA AAAGTGCTCA 8880  
 TCATTGGAAA ACGTTCTTCG GGGCGAAAAC TCTCAAGGAT CTTACCGCTG TTGAGATCCA 8940  
 GTTCGATGTA ACCCACTCGT GCACCCAACT GATCTTCAGC ATCTTTTACT TTCACCAGCG 9000

TTTCTGGGTG AGCAAAAACA GGAAGGCAAA ATGCCGCAAA AAAGGGAATA AGGGCGACAC 9060  
 GGAAATGTTG AATACTCATA CTCTTCCTTT TTCAATATTA TTGAAGCATT TATCAGGGTT 9120  
 ATTGTCTCAT GAGCGGATAC ATATTTGAAT GTATTTAGAA AAATAACAA ATAGGGGTTC 9180  
 CGCGCACATT TCCCCGAAAA GTGCCACCT 9209

(4) INFORMATION FOR SEQ ID NO: 3:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 54 bases
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: DNA (genomic)

(iii) HYPOTHETICAL: yes

(iv) ANTI-SENSE: no

(ix) SEQUENCE DESCRIPTION: SEQ ID NO: 3:

5' ATC ACA GAT CTC TCA CCA TGG ATT TTC AGG TBC AGA TTA TCA GCT 52  
 TC 3' 2

(5) INFORMATION FOR SEQ ID NO: 4:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 30 bases
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: DNA (genomic)

(iii) HYPOTHETICAL: yes

(iv) ANTI-SENSE: yes

(ix) SEQUENCE DESCRIPTION: SEQ ID NO: 4:

5' TGC AGC ATC CGT ACG TTT GAT TTC CAG CTT 3' 30

(6) INFORMATION FOR SEQ ID NO: 5:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 384 bases  
 (B) TYPE: nucleic acid  
 (C) STRANDEDNESS: single  
 (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: DNA (genomic)

(iii) HYPOTHETICAL: yes

(iv) ANTI-SENSE: no

(ix) SEQUENCE DESCRIPTION: SEQ ID NO: 5:

```

ATG GAT TTT CAG GTG CAG ATT ATC AGC TTC CTG CTA ATC AGT GCT TCA GTC   51
ATA ATG TCC AGA GGG CAA ATT GTT CTC TCC CAG TCT CCA GCA ATC CTG TCT   102
GCA TCT CCA GGG GAG AAG GTC ACA ATG ACT TGC AGG GCC AGC TCA AGT GTA   153
AGT TAC ATC CAC TGG TTC CAG CAG AAG CCA GGA TCC TCC CCC AAA CCC TGG   204
ATT TAT GCC ACA TCC AAC CTG GCT TCT GGA GTC CCT GTT CGC TTC AGT GGC   255
AGT GGG TCT GGG ACT TCT TAC TCT CTC ACA ATC AGC AGA GTG GAG GCT GAA   306
GAT GCT GCC ACT TAT TAC TGC CAG CAG TGG ACT AGT AAC CCA CCC ACG TTC   357
GGA GGG GGG ACC AAG CTG GAA ATC AAA                                   384

```

(7) INFORMATION FOR SEQ ID NO: 6:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 27 bases  
 (B) TYPE: nucleic acid  
 (C) STRANDEDNESS: single  
 (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: DNA (genomic)

(iii) HYPOTHETICAL: yes

(iv) ANTI-SENSE: no

(ix) SEQUENCE DESCRIPTION: SEQ ID NO: 6:

5' GCG GCT CCC ACG CGT GTC CTG TCC CAG 3'

27

## (8) INFORMATION FOR SEQ ID NO: 7:

## (i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 29 bases  
 (B) TYPE: nucleic acid  
 (C) STRANDEDNESS: single  
 (D) TOPOLOGY: linear

## (ii) MOLECULE TYPE: DNA (genomic)

## (iii) HYPOTHETICAL: yes

## (iv) ANTI-SENSE: yes

## (ix) SEQUENCE DESCRIPTION: SEQ ID NO: 7:

5' GGS TGT TGT GCT AGC TGM RGA GAC RGT GA 3' 29

## (9) INFORMATION FOR SEQ ID NO: 8:

## (i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 420 bases  
 (B) TYPE: nucleic acid  
 (C) STRANDEDNESS: single  
 (D) TOPOLOGY: linear

## (ii) MOLECULE TYPE: DNA (genomic)

## (iii) HYPOTHETICAL: yes

## (iv) ANTI-SENSE: no

## (ix) SEQUENCE DESCRIPTION: SEQ ID NO: 8:

ATG GGT TGG AGC CTC ATC TTG CTC TTC CTT GTC GCT GTT GCT ACG CGT GTC 51  
 CTG TCC CAG GTA CAA CTG CAG CAG CCT GGG GCT GAG CTG GTG AAG CCT GGG 102  
 GCC TCA GTG AAG ATG TCC TGC AAG GCT TCT GGC TAC ACA TTT ACC AGT TAC 153  
 AAT ATG CAC TGG GTA AAA CAG ACA CCT GGT CGG GGC CTG GAA TGG ATT GGA 204  
 GCT ATT TAT CCC GGA AAT GGT GAT ACT TCC TAC AAT CAG AAG TTC AAA GGC 255  
 AAG GCC ACA TTG ACT GCA GAC AAA TCC TCC AGC ACA GCC TAC ATG CAG CTC 306  
 AGC AGC CTG ACA TCT GAG GAC TCT GCG GTC TAT TAC TGT GCA AGA TCG ACT 357  
 TAC TAC GGC GGT GAC TGG TAC TTC AAT GTC TGG GGC GCA GGG ACC ACG GTC 408  
 ACC GTC TCT GCA 420

## Claims

1. An immunologically active, chimeric anti-CD20 antibody for use as a medicament.

2. An immunologically active, chimeric anti-CD20 antibody for use as a medicament in the treatment of B cell lymphoma in a human.
- 5 3. An immunologically active, chimeric anti-CD20 antibody produced from a transfectoma comprising anti-CD20 in TCAE 8 (within ATCC deposit number 69119).
4. A monoclonal antibody obtainable from a hybridoma which secretes anti-CD20 antibody, said hybridoma being identified by American Type Culture Collection deposit number HB 11388.
- 10 5. An antibody according to claim 3 or claim 4, for use as a medicament.
6. An antibody according to any one of the preceding claims which is radiolabelled.
- 15 7. An antibody according to claim 6, which is radiolabelled with a radiolabel selected from yttrium (90), indium (111), and iodine (131).
8. An antibody according to claim 7, wherein the radiolabel is yttrium (90).
- 20 9. Use of an antibody according to any one of the preceding claims for the manufacture of a medicament for the treatment of B cell lymphoma in a human.
10. Use according to claim 9, wherein the amount of said antibody administered to said human is between about 0.001 to about 30 milligrams of antibody per kilogram body weight of said human ("mg/kg").
- 25 11. Use according to claim 9 or claim 10, wherein a second therapeutically effective amount of at least one immunologically active, chimeric anti-CD20 antibody is administered to said human.
- 30 12. Use according to claim 11, wherein said second therapeutically effective amount of said antibody is administered to said human within about seven days of first administration of said antibody to said human.
- 35 13. Use according to claim 11 or claim 12, wherein a third therapeutically effective amount of at least one immunologically active chimeric anti-CD20 antibody is administered to said human.
14. Use according to claim 13, wherein said third therapeutically effective amount of said antibody is administered to said human within about fourteen days of first administration of said antibody to said human.
15. Use according to claim 9 or claim 10, wherein a radiolabelled anti-CD20 antibody is administered to said human at a second administration period.
- 40 16. Use according to claim 15, wherein the radiolabelled anti-CD20 antibody is obtainable from a hybridoma which secretes anti-CD20 antibody, said hybridoma being identified by American Type Culture Collection deposit number HB 11388.
- 45 17. A hybridoma which secretes anti-CD20 antibody, said hybridoma being identified by American Type Culture Collection deposit number HB 11388.






FIG. 1

LINKER #1 15bp | SV40 ORIGIN=332bp  
 GACGTGCGGG CCGCTCTAGG CCTCCAAAAA AGCCTCCTCA CTACTTCTGG AATAGCTCAG 60  
 AGGCCGAGGC GGCCTCGGCC TCTGCATAAA TAAAAAAAT TAGTCAGCLA TGCATGGGGC 120  
 GGAGAATGGG CGGAAGTGGG CGGAGTTAGG GGCGGGATGG GCGGAGTTAG GGGCGGGACT 180  
 ATGGTTGCTG ACTAATTGAG ATGCATGCTT TGCATACTTC TGCTGCTGG GGAGCCTGGG 240  
 GACTTTCCAC ACCTGGTTGC TGAATAATTG AGATGCATGC TTTGCATACT TCTGCCTGCT 300  
 GGGGAGCCTG GGGACTTTCC ACACCCTAAC TGACACACAT TCCACAGCAAT TAATTCCCCT 360  
 AGTTATTAAT AGTAATCAAT TACGGGGTCA TTAGTTTATA GCCCATATAT GGAGTTCCGC 420  
 GTTACATAAC TTACGGTAAA TGGCCCGCCT GGCTGACCGC CCAACGACCC CCGCCATTG 480  
 CMV PROMOTER-ENHANCER=567bp  
 ACGTCAATAA TGACGTATGT TCCCATAGTA ACGCCAATAG GGACTTTCCA TTGACGTCAA 540  
 TGGGTGGACT ATTTACGGTA AACTGCCCCAC TTGGCAGTAC ATCAAGTGTA TCATATGCCA 600  
 AGTACGCCCC CTATTGACGT CAATGACGGT AAATGGCCCG CCTGGCATTG TGCCAGTAC 660  
 ATGACCTTAT GGGACTTTCC TACTTGGCAG TACATCTACG TATTAGTCAT CGCTATTACC 720  
 ATGGTGATGC GGTTTTGGCA GTACATCAAT GGGCGTGGAT AGCGGTTTGA CTCACGGGGA 780  
 TTTCCAAGTC TCCACCCCAT TGACGTCAAT GGGAGTTTGT TTTGGCACCA AAATCAACGG 840  
 GACTTTCCAA AATGTCGTAA CAACTCCGCC CCATTGACGC AAATGGGCGG TAGGCGTGTA 900  
 CGGTGGGAGG TCTATATAAG CAGAGCTGGG TACGTGAACC GTCAGATCGC CTGGAGACGC 960  
 Bgl II | LINKER #3=76bp | LEADER=60bp  
 CATCACAGAT CTCTACCAT GAGGGTCCCC GCTCAGCTCC TGGGGCTCCT GCTGCTCTGG 1020  
 CTCCCAGGTG CACGATGTGA TGGTACCAAG GTGGAAATCA AACGTACGGT GGCTGCACCA 1080  
 TCTGTCTTCA TCTTCCCGCC ATCTGATGAG CAGTTGAAAT CTGGAAGTGC CTCTGTTGTG 1140  
 TGCTGCTGA ATAATTCTA TCCAGAGAG GCCAAAGTAC AGTGAAGGT GGATAACGCC 1200  
 HUMAN KAPPA CONSTANT 324bp 107 AMINO ACID & STOP CODON  
 CTCCAATCGG GTAACCTCCA GGAGAGTGTG ACAGAGCAGG ACAGCAAGGA CAGCACCTAC 1260  
 AGCCTCAGCA GCACCCTGAC GCTGAGCAAA GCAGACTACG AGAAACACAA AGTCTACGCC 1320  
 TGCGAAGTCA CCCATCAGGG CCTGAGCTCG CCCGTCACAA AGAGCTTCAA CAGGGGAGAG 1380  
 STOP  
 LIGHT  
 CHAIN | Eco RI | LINKER #4=85bp  
 TGTGAAATTC AGATCCGTTA ACGGTTACCA ACTACCTAGA CTGGATTCTG GACAACATGC 1440  
 GGCCGTGATA TCTACGTATG ATCAGCCTCG ACTGTGCCTT CTAGTTGCCA GCCATCTGTT 1500

FIG. 2A

GTTTGCCCCCT CCCCCGTGCC TTCCTTGACC CTGGAAGGTG CCACTCCCAC TGTCCTTTCC 1560  
 TAATAAAATG AGGAAATTGC ATCGCATTGT CTGAGTAGGT GTCATTCTAT TCTGGGGGGT 1620  
 GGGGTGGGGC AGGACAGCAA GGGGGAGGAT TGGGAAGACA ATAGCAGGCA TGCTGGGGAT 1680  
 GCGGTGGGCT CTATGGAACC AGCTGGGGCT CGACAGCTAT GCCAAGTACS CCCCCTATTG 1740  
 ACGTCAATGA CGGTAAATGG CCCGCCTGGC ATTATGCCCC GTACATGACC TTATGGGACT 1800  
 TTCCTACTTG GCAGTACATC TACGTATTAG TCATCGCTAT TACCATGGTG ATGCGGTTTT 1860  
 GGCAGTACAT CAATGGGCGT GGATAGCGGT TTGACTCACG GGGATTTCCTA AGTCTCCACC 1920  
 CCATTGACGT CAATGGGAGT TTGTTTTGGC ACCAAAATCA ACGGGACTTT CAAAATGTC 1980  
 GTAACAATC CGCCCCATTG ACGCAAATGG GCGGTAGGCG TGTACGGTGG GAGGTCTATA 2040  
 TAAGCAGAGC TGGGTACGTC CTCACATTCA GTGATCAGCA CTGAACACAG ACCCGTCGAC 2100  
 2051 2 2058 9 . LINKER #6=7bp Sal I  
 ATGGGTTGGA GCCTCATCTT GCTCTTCTT GTGCTGTTG CTACGGGTGT CGCTAGCACC 2160  
 START HEAVY CHAIN LEADER=51bp Mlu I 2151 2 Nhe I  
 -5 -4 -3 114 115  
 AAGGGCCCAT CGGTCTTCCC CCTGGCACCC TCCTCCAAGA GCACCTCTGG GGGCACAGCG 2220  
 GCCCTGGGCT GCCTGGTCAA GGACTACTTC CCCGAACCGG TGACGGTGTC GTGGAACCTA 2280  
 GCGGCCCTGA CCAGCGGCGT GCACACCTTC CCGGCTGTCC TACAGTCCIC AGGACTCTAC 2340  
 TCCCTCAGCA GCGTGGTGAC CGTGCCCTCC AGCAGCTTGG GCACCCAGAC CTACATCTGC 2400  
 AACGTGAATC ACAAGCCCAAG CAACACCAAG GTGGACAAGA AAGCAGAGCC CAAATCTTGT 2460  
 GACAAAATC ACACATGCCC ACCGTGCCCA GCACCTGAAC TCCTGGGGGG ACCGTCAGTC 2520  
 TTCCTCTTCC CCCCCAAACC CAAGGACACC CTCATGATCT CCCGGACCCC TGAGGTCACA 2580  
 TGGTGGTGG TGGACGTGAG CCACGAAGAC CCTGAGGTCA AGTTCAACTG GTACGTGGAC 2640  
 GCGTGGAGG TGCATAATGC CAAGACAAAG CCGCGGGAGG AGCAGTACAA CAGCACGTAC 2700  
 CGTGTGGTCA GCGTCTCAC CGTCTGCAC CAGGACTGGC TGAATGGCAA GGACTACAAG 2760  
 TGCAAGGTCT CCAACAAAGC CCTCCCAGCC CCCATCGAGA AAACCATCTC CAAAGCCAAA 2820  
 GGGCAGCCCC GAGAACCACA GGTGTACACC CTGCCCCCAT CCCGGGATGA GCTGACCAGG 2880  
 AACCAGGTCA GCCTGACCTG CCTGGTCAAA GGCTTCTATC CCAGCGACAT CGCCGTGGAG 2940  
 TGGGAGAGCA ATGGGCAGCC GGAGAACAAC TACAAGACCA CGCCTCCCGT GCTGGACTCC 3000

FIG. 2B

GACGGCTCCT TCTTCCTCTA CAGCAAGCTC ACCGTGGACA AGAGCAGCTG GCAGCAGGGG 3060  
 AACGTCTTCT CATGCTCCGT GATGCATGAG GCTCTGCACA ACCACTACAC GCAGAAAGAGC 3120  
 STOP HEAVY CHAIN |Bam HI LINKER #7=81bp  
 CTCTCCCTGT CTCCGGGTAA ATGAGGATCC GTTAACGGTT ACCAACTACC TAGACTGGAT 3180  
 3144 15 3225 16  
 TCGTGACAAC ATGCGGGCCGT GATATCTACG TATGATCAGC CTCGACTGTG CTTTCTAGTT 3240  
 GCCAGCCATC TGTTGTTTGC CCTCCCCCG TGCCTTCCTT GACCCTGGAA GGTGCCACTC 3300  
 BOVINE GROWTH HORMONE POLYADENYLATION REGION=231bp  
 CCACTGTCCT TTCCTAATAA AATGAGGAAA TTGCATCGCA TTGTCTGAGT AGGTGTCATT 3360  
 CTATTCTGGG GGGTGGGGTG GGGCAGGACA GCAAGGGGGA GGATTGGGAA GACAATAGCA 3420  
 LINKER #8=34bp  
 GGCATGCTGG GGATGCGGTG GGCTCTATGG AACCAGCTGG GGCTCGACAG CGCTGGATCT 3480  
 3456 17  
 CCGATCCCC | AGCTTTGCTT CTCAATTTCT TATTTGCATA ATGAGAAAAA AAGGAAAAAT 3540  
 3490 1  
 AATTTTAAAC CCAATTCAGT AGTTGATTGA GCAAATGCGT TGCCAAAAAG GATGCTTTAG 3600  
 MOUSE BETA GLOBIN MAJOR PROMOTER=366bp  
 AGACAGTGTT CTCTGCACAG ATAAGGACAA ACATTATTCA GAGGGAGTAC CCAGAGCTGA 3660  
 GACTCCTAAG CCAGTGAGTG GCACAGCATT CTAGGGAGAA ATATGCTTGT CATCACCGAA 3720  
 GCCTGATTCC GTAGAGCCAC ACCTTGGTAA GGGCCAATCT GCTCACACAG GATAGAGAGG 3780  
 GCAGGAGCCA GGGCAGAGCA TATAAGGTGA GGTAGGATCA GTTGCTCCTC ACATTTGCTT 3840  
 LINKER #9=19bp 5' UNTRANSLATED DHFR=82bp  
 CTGACATAGT TGTGTTGGGA GCTTGGATAG CTTGGACAGC TCAGGGCTGC GATTTGCGGC 3900  
 3856 17 3875 16  
 CAAACTTGAC GGCAATCCTA GCGTGAAGGC TGGTAGGATT TTATCCCCGC TGCCATCATG 3960  
 3957 18  
 GTTCGACCAT TGAAGTGCAT CGTCGCCGTG TCCCAAAATA TGGGGATTGG CAAGAACGGA 4020  
 GACCTACCCT GGCCTCCGCT CAGGAACGAG TTCAAGTACT TCCAAAGAAT GACCACAACC 4080  
 TCTTCAGTGG AAGGTAAACA GAATCTGGTG ATTATGGGTA GGAAAACCTG GTTCTCCATT 4140  
 MOUSE DHFR=564bp=187 AMINO ACID & STOP CODON  
 CCTGAGAAGA ATCGACCTTT AAAGGACAGA ATTAATATAG TTCTCAGTAG AGAACTCAA 4200  
 GAACCACCAC GAGGAGCTCA TTTTCTTGCC AAAAGTTTGG ATGATGCCTT AAGACTTATT 4260  
 GAACAACCGG AATTGGCAAG TAAAGTAGAC ATGGTTTGGG TAGTCGGAGG CAGTTCTGTT 4320  
 TACCAGGAAG CCATGAATCA ACCAGGCCAC CTTAGACTCT TTGTGACAAG GATCATGCAG 4380  
 GAATTTGAAA GTGACACGTT TTTCCAGAA ATTGATTTGG GGAAATATAA ACTTGTCCCA 4440  
 GAATACCCAG GCGTCCTCTC TGAGGTCCAG GAGGAAAAAG GCATCAAGTA TAAGTTTGAA 4500

FIG. 2C

STOP DHFR  
 GTCTACGAGA AGAAAGACTA AC-GGAAGAT GCTTTCAAGT TCTCTGCTCC CCTCCTAAAG 4560  
 4521' 2  
 3' UNTRANSLATED DHFR=82bp  
 TCATGCATTT TTATAAGACC ATGGGACTTT TGCTGGCTTT AGATCAGCTT CGACTGTCTT 4620  
 4603' 4 4613' 4  
 TTTAGTTGC CAGCCATCTG TTSTTTGCCC CTCCCCCGTG CCTTCCTTGA CCCTGGAAGG 4680  
 BOVINE GROWTH HORMONE POLYADENYLATION REGION=231bp  
 TGGCACTCCC ACTGTCTCTT CTAATAAAA TGAGGAAATT GCATCGCATT GTCTGAGTAG 4740  
 GTGTCATTCT ATTCTGGGGG GTGGGGTGGG GCAGGACAGC AAGGGGGAGG ATTGGGAAGA 4800  
 CAATAGCAGG CATGCTGGGG ATGGGGTGGG CTCTATGGAA CCAGCTGGGG CTGAGCTAC 4860  
 4844' 5  
 TAGCTTTGCT TCTCAATTC TTATTTGCAT AATGAGAAAA AAAGGAAAAT TAATTTTAAC 4920  
 ACCAATTCAG TAGTTGATTG AGCAAAATGCG TTGCCAAAAA GGATGCTTTA GAGACAGTGT 4980  
 MOUSE BETA GLOBIN MAJOR PROMOTER=366bp  
 TCTCTGCACA GATAAGGACA AACATTATTC AGAGGGAGTA CCCAGAGCTG AGACTCCTAA 5040  
 GCCAGTGAGT GGCACAGCAT TCTAGGGAGA AATATGCTTG TCATCACCGA AGCCTGATTC 5100  
 CGTAGAGCCA CACCTTGGTA AGGGCCAAATC TGCTCACACA GGATAGAGAG GGCAGGAGCT 5160  
 AGGGCAGAGC ATATAAGGTG AGGTAGGATC AGTTGCTCCT CACATTGCT TCTGACATAG 5220  
 TTGTGTTGGG AGCTTGGATC GATCCTCTAT GGTGAACAA GATGGATTGC ACGCAGGTTT 5280  
 5227' 8 5248' 9  
 TCCGGCCGCT TGGGTGGAGA GGCTATTCGG CTATGACTGG GCACAACAGA CAATCGGCTG 5340  
 CTCTGATGCC GCCGTGTTCC GGCTGTCAGC GCAGGGGGCG CCGGTTCTTT TTGTCAAGAC 5400  
 NEOMYCIN PHOSPHOTRANSFERASE  
 CGACCTGTCC GGTGCCCTGA ATGAAGTGA GGACGAGGCA GCGCGGCTAT CGTGGCTGGC 5460  
 795bp=264 AMINO ACIDS & STOP CODON  
 CACGACGGGC GTTCCTTGCG CAGCTGTGCT CGACGTTGTC ACTGAAGCGG GAAGGGACTG 5520  
 GCTGCTATTG GGCGAAGTGC CGGGGCAGGA TCTCCTGTCA TCTACCTTG CTCCTGCCGA 5580  
 GAAAGTATCC ATCATGGCTG ATGCAATGCG GCGGCTGCAT ACGCTTGATC CGGCTACCTG 5640  
 CCCATTCGAC CACCAAGCGA AACATCGCAT CGAGCGAGCA CGTACTCGGA TGAAGCCGG 5700  
 TCTTGTGGAT CAGGATGATC TGGACGAAGA GCATCAGGGG CTCGCGCCAG CCGAACTGTT 5760  
 CGCCAGGCTC AAGGCGCGCA TGCCCGACGG CGAGGATCTC GTCGTGACCC ATGGCGATGC 5820  
 CTGCTTGCCG AATATCATGG TGGAAAATGG CCGCTTTTCT GGATTCATCG ACTGTGGCCG 5880  
 GCTGGGTGTG GCGGACCGCT ATCAGGACAT AGCGTTGGCT ACCCGTGATA TTGCTGAAGA 5940  
 GCTTGGCGGC GAATGGGCTG ACCGCTTCCT CGTGCTTTAC GGTATCGCCG CTTCCCGATTG 6000

FIG. 2D

GCAGCGCATC GCCTTCTATC GCCTTCTTGA CGAGTTCTTC <sup>STOP NEOI</sup> TCAGCGGGAC TCTGGCGTTC 6060  
 GAAATGACCG ACCAAGCGAC GCGCAACCTG CCATCACGAG ATTTCGATTC CACCGCCGCC 6120  
 TTCTATGAAA GGTGCGGCTT CGGAATCGTT TTCCGGGACG CCGGCTGGAT GATCCTCCAG 6180  
 CGCGGGGATC TCATGCTGGA GTTCTTCGCC CACCCCAACT TGTATTATGC AGCTTATAAT 6240  
 GGTTACAAAT AAAGCAATAG CATCACAAAT TTCACAAATA AAGCATTTTT TTCACTGCAT 6300  
 TCTAGTTGTG GTTTGTCCAA ACTCATCAAT CTATCTTATC <sup>SV40 POLY A EARLY=133bp</sup> ATGCTGGAT CGCGGCCGCG 6360  
 ATCCCGTCCA GAGCTTGGCG TAATCATGGT CATAGCTGTT TCCTGTGTGA AATTGTTATC 6420  
 CGCTCACAAT TCCACACAAC ATACGAGCCG GAAGCATAAA GTGTAAAGCC TGGGGTGCCT 6480  
 AATGAGTCAG CTAATCACA TTAATTGCGT TCGCTCACT GCGCGCTTTC CAGTCGGGAA 6540  
 ACCTGTCTG CCAGCTGCAT TAATGAATCG GCCAACGCGC GGGGAGAGGC GGTTTGCGTA 6600  
 TTGGGCGCTC TTCCGCTTCC TCGCTCACTG <sup>PVC 19</sup> ACTCGCTGCG CTCGGTCGTT CGGCTGCGGC 6660  
 GAGCGGTATC AGCTCACTCA AAGGCGGTAA TACGGTTATC CACAGAATCA GGGGATAACG 6720  
 CAGGAAAGAA CATGTGAGCA AAAGGCCAGC AAAAGGCCAG GAACCGTAAA AAGGCCGCT 6780  
 TGCTGGCGTT <sup>6792=BACTERIAL ORIGIN OF REPLICATION</sup> TTTCATAGG CTCGCGCCCC CTGACGAGCA TCACAAAAAT CGACGCTCAA 6840  
 GTCAGAGGTG GCGAAACCCG ACAGGACTAT AAAGATACCA GCGGTTTCCC CCTGGAAGCT 6900  
 CCCTCGTGCG CTCTCCTGTT CCGACCCTGC CGCTTACCGG ATACCTGTCC GCCTTTCTCC 6960  
 CTTCGGGAAG CGTGGCGCTT TCTCAATGCT CACGCTGTAG GTATCTCAGT TCGGTGTAGG 7020  
 TCGTTGCTC CAAGCTGGGC TGTGTGCACG AACCCCCCGT TCAGCCCGAC CGCTGCGCCT 7080  
 TATCCGGTAA CTATCGTCTT GAGTCCAACC CGGTAAGACA CGACTTATCG CCACTGECAG 7140  
 CAGCCACTGG TAACAGGATT AGCAGAGCGA GGTATGTAGG CGGTGCTACA GAGTTCTTGA 7200  
 AGTGGTGGCC TAACTACGGC TACACTAGAA GGACAGTATT TGGTATCTGC GCTCTGCTGA 7260  
 AGCCAGTTAC CTTCGGAAAA AGAGTTGGTA GCTCTTGATC CGGCAAAACA ACCACCGCTG 7320  
 GTAGCGGTGG TTTTTTTGTT TGCAAGCAGC AGATTACGCG CAGAAAAAAA GGATCTCAAG 7380  
 AAGATCCTTT GATCTTTTCT ACGGGGTCTG ACGCTCAGTG GAACGAAAAC TCACGTTAA 7440  
 GGATTTTGGT CATGAGATTA TCAAAAAGGA TCTTCACCTA GATCCTTTTA AATTAAAAAT 7500

FIG. 2E

GAAGTTTTAA ATCAATCTAA AGTATATATG AGTAAACTTG GTCTGACAGT TACCAATGCT 7550  
 TAATCAGTGA GGCACCTATC TCACCGATCT GTCTATTTCTG TTCATCCATA GTTGCCCTGAC 7620  
 TCCCCGTCGT GTAGATAACT ACGATACGGG AGGGCTTACC ATCTGGCCCC AGTGCTGCAA 7680  
 TGATACCGCG AGACCCACGC TCACCGGCTC CAGATTTATC AGCAATAAAC CAGCCAGCCG 7740  
 GAAGGGCCGA GCGCAGAAGT GGTCTTGCAA CTTTATCCGC CTCCATCCAG TCTATTAAATT 7800  
 GTTGCCGGGA AGCTAGAGTA AGTAGTTCGC CAGTTAATAG TTTGCGCAAC GTTGTTGCCA 7860  
 TTGCTACAGG CATCGTGGTG TCACGCTCGT CGTTTGGTAT GGCTTCATTC AGCTCCGGTT 7920  
 CCCAACGATC AAGGCGAGTT ACATGATCCC CCATGTTGTG CAAAAAAGCG GTTAGCTCCT 7980  
 TCGGTCCTCC GATCGTTGTC AGAAGTAAGT TGGCCGCAGT GTTATCACTC ATGGTTATGG 8040  
 CAGCACTGCA TAATTCTCTT ACTGTCATGC CATCCGTAAG ATGCTTTTCT GTGACTGGTG 8100  
 AGTACTCAAC CAAGTCATTC TGAGAATAGT GTATGCGGCG ACCGAGTTGC TCTTGCCCGG 8160  
 CGTCAATACG GGATAATACC GCGCCACATA GCAGAACTTT AAAAGTGCTC ATCATTGGAA 8220  
 AACGTTCTTC GGGGCGAAAA CTCTCAAGGA TCTTACCGCT GTTGAGATCC AGTTCGATGT 8280  
 AACCCACTCG TGCACCCAAC TGATCTTCAG CATCTTTTAC TTTACCAGC GTTCTGGGT 8340  
 GAGCAAAAAC AGGAAGGCAA AATGCCGCAA AAAAGGGAAT AAGGGCGACA CGGAAATGTT 8400  
 GAATACTCAT ACTCTTCCTT TTTCAATATT ATTGAAGCAT TTATCAGGGT TATTGTCTCA 8460  
 TGAGCGGATA CATATTTGAA TGTATTTAGA AAAATAAACA AATAGGGGTT CCGCGCACAT 8520  
 TTCCCCGAAA AGTGCCACCT

*FIG. 2F*

LINKER #1=15bp  
 GACGTGCGGG CCGCTCTAGG CCTCCAAAA AGCCTCCTCA CTACTTCTGG AATAGCTCAG 60  
 15 6  
 AGGCCGAGGC GGCCTCGGCC TCTGCATAAA TAAAAAAAT TAGTCACCA TGCATGGGG 120  
 SV40 ORIGIN=332bp  
 GGAGAATGGG CGGAAGTGGG CGGAGTTAGG GCGGGGATGG GCGGAGTTAG GGGCGGGACT 180  
 ATGGTTGCTG ACTAATTGAG ATGCATGCTT TGCATACTTC TGCCTGCTGG GGAGCCTGGG 240  
 GACTTTCCAC ACCTGGTTGC TGAATAATTG AGATGCATGC TTTGCATACT TCTGCCTGCT 300  
 GGGGAGCCTG GGGACTTTCC ACACCCTAAC TGACACACAT TCCACAGAAAT TAATTCCCCT 360  
 347 8  
 AGTTATTAAT AGTAATCAAT TACGGGGTCA TTAGTTCATA GCCCATATAT GGAGTTCCGC 420  
 GTTACATAAC TTACGGTAAA TGGCCCGCCT GGCTGACCGC CCAACGAGCC CCGCCCATGG 480  
 ACGTCAATAA TGACGTATGT TCCCATAGTA ACGCCAATAG GGACTTTCCA TTGACGTCAA 540  
 CVM PROMOTER-ENHANCER=567bp  
 TGGGTGGACT ATTTACGGTA AACTGCCAC TTGGCAGTAC ATCAAGTGTA TCATATGCCA 600  
 AGTACGCCCC CTATTGACGT CAATGACGGT AAATGGCCCG CCTGGGATTA TGCACAGTAC 660  
 ATGACCTTAT GGGACTTTCC TACTTGGCAG TACATCTACG TATTAGTCAT CGCTATTACC 720  
 ATGGTGATGC GGTTTTGGCA GTACATCAAT GGGCGTGGAT AGCGGTTTGA CTCACGGGGA 780  
 TTTCCAAGTC TCCACCCCAT TGACGTCAAT GGGAGTTTGT TTTGGCACCA AAATCAACGG 840  
 GACTTTCCAA AATGTCGTAA CAACTCCGCC CCATTGACGC AAATGGGCGG TAGGCGTGTA 900  
 CGGTGGGAGG TCTATATAAG CAGAGCTGGG TACGTGAACC GTCAGATCGC CTGGAGACGC 960  
 927 8 934 5  
 Bgl 2 START LIGHT CHAIN NATURAL LEADER=66bp  
 CATCACAGAT CTCTCACTAT GGATTTTCAG GTGCAGATTA TCAGCTTCCT GCTAATCAGT 1020  
 978 9  
 GCTTCAGTCA TAATGTCCAG AGGACAAATT GTTCTCTCCC AGTCTCCAGC AATCCTGTCT 1080  
 1044 5+1  
 GCATCTCCAG GGGAGAAGGT CACAATGACT TGCAGGGCCA GCTGAAGTGT AAGTTACATC 1140  
 CACTGGTTCC AGCAGAAGCC AGGATCCTCC CCCAAACCCT GGATTTATGC CACATCCAAC 1200  
 LIGHT CHAIN VARIABLE REGION 318bp 106 AMINO ACID  
 CTGGCTTCTG GAGTCCCTGT TCGCTTCAGT GGCAGTGGGT CTGGGACTTC TTA CTCTCTCTC 1260  
 ACCATCAGCA GAGTGGAGGC TGAAGATGCT GCCACTTATT ACTGCCAGCA GTGGACTAGT 1320  
 AACCACCCA CGTTCGGAGG GGGGACCAAG CTGGAAATCA AACGTACGGT GGCTGCACCA 1380  
 1362 3  
 TCTGTCTTCA TCTTCCCGCC ATCTGATGAG CAGTTGAAAT CTGGAACGCG CTCTGTTGTG 1440  
 TGCCTGCTGA ATAATTCTA TCCAGAGAG GCCAAAGTAC AGTGAAGGT GGATAACGCC 1500

FIG. 3A



HUMAN KAPPA CONSTANT=324bp=107 AMINO ACID & STOP CODON  
 CTCCAATCGG GTAACCTCCA GGAGAGTGTC ACAGAGCAGG ACAGCAAGGA CAGCACCTAC 1560  
 AGCCTCAGCA GCACCCTGAC GCTGAGCAAA GCAGACTACG AGAAACACAA AGTCTACGCC 1620  
 TCGGAAGTCA CCCATCAGGG CCTGAGCTCG CCCGTCAAA AGAGCTTCAA CAGGGGAGAG 1680  
 STOP  
 LIGHT  
 CHAIN/Eco RI LINKER #4=81bp  
 TGTTCGAATTC AGATCCGTTA ACGGTTACCA ACTACCTAGA CTGGATTCGT GACAACATGC 1740  
 1646 17  
 GGCCGTGATA TCTACGTATG ATCAGCCTCG ACTGTGCCTT CTAGTTGCCA GCCATCTGTT 1800  
 1771 12  
 GTTTGCCCCCT CCCCCGTGCC TTCCTTGACC CTGGAAGGTG CCACTCCCAC TGTCCTTTCC 1860  
 TAATAAAATG AGGAAATTGC ATCGCATTGT CTGAGTAGGT GTCATTCTAT TCTGGGGGGT 1920  
 BOVINE GROWTH HORMONE POLYADENYLATION REGION=231bp  
 GGGGTGGGGC AGGACAGCAA GGGGGAGGAT TGGGAAGACA ATAGCAGGCA TGCTGGGGAT 1980  
 LINKER #5=15bp  
 GCGGTGGGCT CTATGGAACC ACTGGGGCT CGACGCTAT GCCAAGTACG CCCCTATTG 2040  
 2002 3 2017 8  
 ACGTCAATGA CGGTAAATGG CCCGCCTGGC ATTATGCCCA GTACATGACC TTATGGGACT 2100  
 TTCCTACTTG GCAGTACATC TACGTATTAG TCATCGCTAT TACCATGGTG ATGCGGTTTT 2160  
 CMV PROMOTER-ENHANCER=334bp  
 GGCAGTACAT CAATGGGCGT GGATAGCGGT TTGACTCAGG GGGATTTCCA AGTCTCCACC 2220  
 CCATTGACGT CAATGGGAGT TTGTTTTGGC ACCAAAATCA ACGGGACTTT CCAAAATGTC 2280  
 GTAACAATC CGCCCCATTG ACGCAAATGG GCGGTAGGCG TGTACGGTGG GAGGTCTATA 2340  
 LINKER #6=7bp Sal I  
 TAAGCAGAGC TGGGTACGTC CTCACATTCA GTGATCAGCA CTGAACACAG ACCCGTCCAC 2400  
 2351 2 2358 9  
 START  
 HEAVY CHAIN SYNTHETIC & NATURAL LEADER Mlu I 2457 8  
 ATGGGTGGA GCCTCATCTT GCTCTTCCTT GTCGCTGTG CTACGCGTGT CCTGTCCAG 2460  
 2401 -5 -4 -3 -2 -1 +1  
 GTACAACTGC AGCAGCCTGG GGCTGAGCTG GTGAAGCCTG GGGCCTCAGT GAAGATGTCC 2520  
 TGCAAGGCTT CTGGCTACAC ATTTACCACT TACAATATGC ACTGGGTAAA ACAGACACCT 2580  
 HEAVY CHAIN VARIABLE=363bp=121 AMINO ACID  
 GGTGGGGGCC TGAATGGAT TGGAGCTATT TATCCCGGAA ATGGTGATAC TTCCTACAAT 2640  
 CAGAAGTTCA AAGGCAAGGC CACATTGACT GCAGACAAAT CCTCCAGCAC AGCCTACATG 2700  
 CAGCTCAGCA GCCTGACATC TGAGGACTCT GCGGTCTATT ACTGTGCAAG ATCGACTTAC 2760  
 TACGGCGGTG ACTGGTACTT CAATGTCTGG GGCGCAGGGA CCACGGTCAC CGTCTCTGCA 2820  
 Nhe I  
 GCTAGCACCA AGGGCCCATC GGTCTTCCCC CTGGCACCTT CCTCCAAGAG CACCTCTGGG 2880  
 GGCACAGCGG CCCTGGGCTG CCTGGTCAAG GACTACTTCC CCGAACCGLT GACGGTGTGG 2940  
 HUMAN GAMMA 1 CONSTANT=993bp  
 TGGAACTCAG GCGCCCTGAC CAGCGGCGTG CACACCTTCC CGGCTGTCCT ACAGTCCTCA 3000

FIG. 3B

330 AMINO ACID & STOP CODON  
 GGACTCTACT CCCTCAGCAG CGTGGTGACC G-GCCCTCCA GCAGCTTGGG CACCCAGACC 3060  
 TACATCTGCA ACGTGAATCA CAAGCCCAGC AACACCAAGG TGGACAAAGAA AGCAGAGCCC 3120  
 AAATCTTGTG AAAAACTCA CACATGCCCA CCGTGCCAG CACCTGAACT CCTGGGGGGA 3180  
 CCGTCAGTCT TCCTCTTCCC CCCAAAACCC AAGGACACCC TCATGATCTC CCGGACCCCT 3240  
 GAGGTCACAT GCGTGGTGGT GGACGTGAGC CACGAAGACC CTGAGGTCAA GTTCAACTGG 3300  
 TACGTGGACG GCGTGGAGGT GCATAATGCC AAGACAAAGC CGCGGGAGGA GCAGTACAAC 3360  
 AGCACGTACC GTGTGGTCAG CGTCCTCACC GTCTGCACC AGGACTGGCT GAATGGCAAG 3420  
 GAGTACAAGT GCAAGGTCTC CAACAAAGCC CTCCCAGCCC CCATCGAGAA AACCATCTCC 3480  
 AAAGCCAAAG GGCAGCCCCG AGAACCACAG GTGTACACCC TGCCCCCATC CCGGGATGAG 3540  
 CTGACCAAGA ACCAGGTCAG CCTGACCTGC CTGGTCAAAG GCTTCTATCC CAGCGACATC 3600  
 GCCGTGGAGT GGGAGAGCAA TGGGCAGCCG GAGAACAAC ACAAGACCAC GCCTCCCGTG 3660  
 CTGGACTCCG ACGGCTCCTT CTTCTCTAC AGCAAGCTCA CCGTGGACAA GAGCAGGTGG 3720  
 CAGCAGGGGA ACGTCTTCTC ATGCTCCGTG ATGCATGAGG CTCTGCACAA CCACTACAGG 3780  
 CAGAAGAGCC TCTCCCTGTC TCCGGGTAAA **STOP HEAVY CHAIN I<sub>H</sub>am HI** **LINKER #7=81bp**  
 3813<sup>4</sup> TCAGGATCCG TTAACGGTTA CCAACTACCT 3840  
 AGACTGGATT CGTGACAACA TGGCGCCGTG ATATCTACGT ATGATCAGCC TCGACTGTGC 3900  
 3894<sup>5</sup>  
 CTTCTAGTTG CCAGCCATCT GTTGTITGCC CCTCCCCCGT GCCTTCCTTG ACCCTGGAAG 3960  
 GTGCCACTCC CACTGTCCTT TCCTAATAAA ATGAGGAAAT TGCATCGCAT TGTCTGAGTA 4020  
**BOVINE GROWTH HORMONE POLYADENYLATION REGION=231bp**  
 GGTGTCAATC TATTCTGGGG GGTGGGGTGG GGCAGGACAG CAAGGGGGAG GATTGGGAAG 4080  
 ACAATAGCAG GCATGCTGGG GATGCGGTGG GCTCTATGGA ACCAGCTGGG GCTCGACAGC 4140  
 4125<sup>6</sup>  
 GCTGGATCTC CCGATCCCA GCTTTGCTTC TCAATTTCTT ATTTGCATAA TGAGAAAAAA 4200  
 AGGAAAATTA ATTTTAACAC CAATTCAGTA GTTGATTGAG CAAATGCGTT GCCAAAAAGG 4260  
**MOUSE BETA GLOBIN MAJOR PROMOTER=386bp**  
 ATGCTTTAGA GACAGTGGTC TCTGCACAGA TAAGGACAAA CATTATTCAG AGGGAGTACC 4320  
 CAGAGCTGAG ACTCCTAAGC CAGTGAGTGG CACAGCATTC TAGGGAGAAA TATGCTTGTC 4380  
 ATCACCGAAG CCTGATTCCG TAGAGCCACA CCTTGGTAAG GGCCAATCTG CTCACACAGG 4440  
 ATAGAGAGGG CAGGAGCCAG GGCAGAGCAT ATAAGGTGAG GTAGGATCAG TTGCTCCTCA 4500

FIG. 3C

CATTGCTTC TGACATAGTT LINKER #9=19bp 5' UNTRANSLATED DHFR=82bp  
 GTGTGGGAG CTTGGATAG 4525'6 TTGGACAGCT CAGCGCTGGC 4544'5 4560  
 ATTTGGGGCC AACTTGACG GCAATCCTAG CGTGAAGGC GGTAGGATTT TATCCCGCT 4620  
START DHFR  
 GCCATC 4626'7 TTGACCATT GAACTGCATC GTCGCCGTGT CCCAAAATAT GGGGATTGGC 4680  
 AAGAACGGAG ACCTACCCTG GCCTCCGCTC AGGAACGAGT TCAAGTACTT CCAAAGAATG 4740  
 ACCACAACCT CTTCACTGGA AGGTAAACAG AATCTGGTGA TTATGGGTAG GAAAACCTGG 4800  
DHFR=564bp=187 AMINO ACID & STOP CODON  
 TTCTCCATTC CTGAGAAGAA TCGACCTTTA AAGGACAGAA TTAATATAGT TCTCAGTAGA 4860  
 GAACTCAAAG AACCACCACG AGGAGCTCAT TTTCTTGCCA AAAGTTTGGG TGATGCCTTA 4920  
 AGACTTATTG AACAACCGGA ATTGGCAAGT AAAGTAGACA TGGTTTGGAT AGTCGGAGGC 4980  
 AGTTCTGTTT ACCAGGAAGC CATGAATCAA CCAGGCCACC TTAGACTCTT TGTGACAAGG 5040  
 ATCATGCAGG AATTTGAAAG TGACACGTTT TTCCCAGAAA TTGATTTGGG GAAATATAAA 5100  
 CTTCTCCCAG AATACCCAGG CGTCCTCTCT GAGGTCCAGG AGGAAAAAGG CATCAAGTAT 5160  
STOP DHFR 3' UNTRANSLATED DHFR=82bp  
 AAGTTTGAAG TCTACGAGAA GAAAGACTAA 5140'1 CAGGAAGATG CTTTCAAGTT CTCTGCTCCC 5220  
LINKER #10  
 CTCCTAAAGC TATGCATTTT TATAAGACCA TGGGACTTTT GCTGGCTTTA GATCAGCCTC 5272'3 5280  
=10bp  
 GACTGTGCCT TCTAGTTGCC AGCCATCTGT TGTTTGCCCC TCCCCCGTGC CTTCTTGAC 5340  
BOVINE GROWTH HORMONE POLYADENYLATION=231bp  
 CCTGGAAGGT GCCACTCCCA CTGTCCTTTC CTAATAAAAT GAGGAAATTG CATCGCATTG 5400  
 TCTGAGTAGG TGTCATTCTA TTCTGGGGGG TGGGGTGGGG CAGGACAGCA AGGGGGAGGA 5460  
LINKER #11  
 TTGGGAAGAC AATAGCAGGC ATGCTGGGGA TCGGGTGGGC TCTATGGAAC CAGCTGGGGC 5513'4 5520  
=17bp  
 TCGAGCTACT AGCTTTGCTT CTCAATTTCT TATTTGCATA ATGAGAAAAA AAGGAAAATT 5580  
5530'1  
 AATTTTAACA CCAATTCAGT AGTTGATTGA GCAAATGCGT TGCCAAAAAG GATGCTTTAG 5640  
MOUSE BETA GLOBIN MAJOR PROMOTER=366bp  
 AGACAGTGTT CTCTGCACAG ATAAGGACAA CTAGGGAGAA ATATGCTTGT CATCACCGAA 5700  
 GACTCCTAAG CCAGTGAGTG GCACAGCATT CTAGGGAGAA ATATGCTTGT CATCACCGAA 5760  
 GCCTGATTCC GTAGAGCCAC ACCTTGGTAA GGGCCAATCT GCTCACACAG GATAGAGAGC 5820  
 GCAGGAGCCA GGGCAGAGCA TATAAGGTGA GGTAGGATCA GTTGCTCCTC ACATTTGCTT 5880  
LINKER #12=21bp START NEO  
 CTGACATAGT TGTGTTGGGA GCTTGGATCG ATCCTCTATG 5896'7 5917'8 GTTGAACAAG ATGGATTGCA 5940  
 CGCAGGTTCT CCGGCCGCTT GGGTGGAGAG GCTATTCGGC TATGACTGGG CACAACAGAC 6000

FIG. 3D

AATCGGCTGC TCTGATGCCG CCGTGTTCCG GCTGTCAGCG CAGGGGCGCC CGGTTCTTTT 6060  
 NEOMYCIN PHOSPHOTRANSFERASE=795bp=264 AMINO ACID & STOP CODON  
 TGTCAAGACC GACCTGTCCG GTGCCCTGAA TGAAGTGCAG GACGAGGCAG CGCGGCTATC 6120  
 GTGGCTGGCC ACGACGGGCG TTCCTTGCGC AGCTGTGCTC GACGTTGTCA CTGAAGCGCC 6180  
 AAGGGACTGG CTGCTATTGG GCGAAGTGCC GGGGCAGGAT CTCCTGTCAT CTCACCTTGC 6240  
 TCCTGCCGAG AAAGTATCCA TCATGGCTGA TGCAATGCGG CGGCTGCATA CGCTTGATCC 6300  
 GGCTACCTGC CCATTCGACC ACCAAGCGAA ACATCGCATC GAGCGAGCAC GTACTCGGAT 6360  
 GGAAGCCGGT CTTGTGATC AGGATGATCT GGACGAAGAG CATCAGGGGC TCGCGCCAGC 6420  
 CGAACTGTTC GCCAGGCTCA AGGCGCGCAT GCCCGACGGC GAGGATCTCG TCGTGACCCA 6480  
 TGGCGATGCC TGCTTGCCGA ATATCATGGT GGAAAATGGC CGCTTTTCTG GATTCATCGA 6540  
 CTGTGGCCGG CTGGGTGTGG CGGACCGCTA TCAGGACATA GCGTTGGCTA CCCGTGATAT 6600  
 TGCTGAAGAG CTTGGCGGCG AATGGGCTGA CCGCTTCCTC GTGCTTTACG GTATCGCCGC 6660  
 TCCCGATTCT CAGCGCATCG CTTCTATCG CTTCTTGAC GAGTTCTTCT ~~GAGCGGGACT~~ 6720  
 STOP NEO  
 6712 3  
 GTGGGGTTCT AAATGACCGA CCAAGCGAGC CCCAACCTGC CATCAGGAGA TTTCGATTCT 6780  
 3' UNTRANSLATED NEO=173bp  
 ACCGCCGCCT TCTATGAAAG GTTGGGCTTC GGAATCGTTT TCCGGGACGC CGGCTGGATG 6840  
 ATCCTCCAGC GCGGGGATCT CATGCTGGAG TTCTTCGCC ACCCAACTT GTTTATTGCA 6900  
 6885 6  
 GCTTATAATG GTTACAAATA AAGCAATAGC ATCACAATTT TCACAAATAA AGCATTTTTT 6960  
 SV40 EARLY POLYADENYLATION REGION=133bp  
 TCACTGCATT CTAGTTGTGG TTTGTCCAAA CTCATCAATC TATCTTATCA TGTCTGGATC 7020  
 7018 9  
 LINKER #13=19bp  
 GCGGCCGCGA TCCCGTCTGAG AGCTTGCGCT AATCATGGTC ATAGCTGTTT CCTGTGTGAA 7080  
 7037 8  
 PUC 19  
 ATTGTTATCC GCTCACAATT CCACACAACA TACGAGCCGG AAGCATAAAG TGTAAGCCT 7140  
 GGGGTGCCTA ATGAGTGAGC TAACTCACAT TAATTGCGTT GCGCTCACTG CCCGCTTTCT 7200  
 AGTCGGGAAA CCTGTGCTGC CAGCTGCATT AATGAATCGG CCAACGCGCG GGGAGAGGCG 7260  
 GTTTGCGTAT TGGGCGCTCT TCCGCTTCCT CGCTCACTGA CTCGCTGCGC TCGGTCGTTT 7320  
 GGCTGCGGCG AGCGGTATCA GCTCACTCAA AGGCGGTAAT ACGGTTATCC ACAGAATCAG 7380  
 GGGATAACGC AGGAAAGAAC ATGTGAGCAA AAGGCCAGCA AAAGGCCAGG AACCGTAAAA 7440  
 7461=BACTERIAL ORIGIN OF REPLICATION  
 AGGCCGCGTT GCTGGCGTTT TCCATAGGC TCCGCCCCCT TGACGAGCAT CACAAAAATC 7500

FIG. 3E

GACGCTCAAG TCAGAGGTGG CGA--ACCGG- CAGGACTATA AAGATACCAG GCGTTTCCCC 7560  
 CTGGAAGCTC CCTCGTGCGC TCTCCTGTTG CGACCCCTGCC GCTTACCGGA TACCTGTCCG 7620  
 CCTTTCTCCC TTCCGGGAAGC GTGCGCGCTTT CTCAATGCTC ACGCTGTAGG TATCTCAGTT 7680  
 CCGTGTAGGT CGTTCCGCTCC AAGCTGGGCT GTGTGCACGA ACCCCCCGTT CAGCCCCGACC 7740  
 GCTGCGCCTT ATCCGGTAAC TATCGTCTTG AGTCCAACCC GGTAAGACAC GACTTATCGC 7800  
 CACTGGCAGC AGCCACTGGT AACAGGATTA GCAGAGCGAG GTATGTAGGC GGTGCTACAG 7860  
 AGTTCTTGAA GTGGTGGCCT AACTACGGCT AACTAGAAG GACAGTATTT GGTATCTCCG 7920  
 CTCTGCTGAA GCCAGTTACC TTCGGAAAAA GAGTTGGTAG CTCTTGATCC GGCAACAAA 7980  
 CCACCGCTGG TAGCGGTGGT TTTTTTGTTC GCAAGCAGCA GATTACGCGC AGAAAAAAG 8040  
 GATCTCAAGA AGATCCTTTG ATCTTTTCTA CGGGGTCTGA CGCTCAGTGG AACGAAACT 8100  
 CACGTAAAGG GATTTTGGTC ATGAGATTAT CAAAAAGGAT CTTACCTAG ATCCTTTTAA 8160  
 ATTAAAAATG AAGTTTTAAA TCAATCTAAA GTATATATGA GTAAACTTGG TCTGACAGTT STOP 8220  
BETA LACTAMASE  
ACCAATGCTT AATCAGTGAG GCACCTATCT CAGCGATCTG TCTATTTCTG TCATCCATAG 8280  
 TTGECTGACT CCCCCTCGTG TAGATACTA CUAACGGGA GGGCTTACCA TCTGGCCCCA 8340  
 GTGCTGCAAT GATACCGCGA GACCCACGCT CACCGGCTCC AGATTTATCA GCAATAAACC 8400  
 BETA LACTAMASE=861bp=286 AMINO ACID & STOP CODON  
 AGCCAGCCGG AAGGGCCGAG CGCAGAAGTG GTCCTGCAAC TTTATCCGCC TCCATCCAGT 8460  
 CTATTAATTG TTGCCGGGAA GCTAGAGTAA GTAGTTCGCC AGTTAATAGT TTGCGCAACG 8520  
 TTGTTGCCAT TGCTACAGGC ATCGTGGTGT CACGCTCGTC GTTTGGTATG GCTTCATTCA 8580  
 GCTCCGGTTC CCAACGATCA AGGCGAGTGA CATGATCCCC CATGTTGTGC AAAAAAGCGG 8640  
 TTAGCTCCTT CGGTCTCTCG ATCGTTGTCA GAAGTAAGTT GGCCGCACTG TTATCACTCA 8700  
 TGGTTATGGC AGCACTGCAT AATTCTCTTA CTGTCATGCC ATCCGTAAGA TGCTTTTCTG 8760  
 TGA CTGGTGA GTACTCAACC AAGTCATTCT GAGAATAGTG TATGCGGCGA CCGAGTTGCT 8820  
 CTTGCCCGGC GTCAATACGG GATAATACCG CGCCACATAG CAGAACTTTA AAAGTGCTCA 8880  
 TCATTGAAA ACGTTCTTCG GGGCGAAAAC TCTCAAGGAT CTTACCGCTG TTGAGATCCA 8940  
 GGTGATGTA ACCCACTCGT GCACCAACT GATCTTCAGC ATCTTTTACT TTCACCAGCG 9000  
 TTTCTGGGTG AGCAAAAACA GGAAGGCAAA ATGCCGCAAA AAAGGGAATA AGGGCGACAC 9060  
 GGAAATGTTG AATACTCATA START BETA LACTAMASE CTCTTCCTTT TTCAATATTA TTGAAGCATT TATCAGGGTT 9120  
 ATTGTCTCAT GAGCGGATAC ATATTGAAT GTATTTAGAA AAATAAACAA ATAGGGGTTT 9180  
 CGCGCACATT TCCCCGAAAA GTGCCACCT

FIG. 3F

FIG. 4

LEADER

-19		-15		-10		-5										
Met	Gly	Trp	Ser	Leu	Ile	Leu	Leu	Phe	Leu	Val	Ala	Val	Ala	Thr	Arg	Val
ATG	GGT	TGG	AGC	CTC	ATC	TTG	CTC	TTC	CTT	GTC	GCT	GTT	GCT	ACG	CGT	GTC
	2409			2418				2427			2436			2445		

-1	+1	FR1		10		15												
Leu	Ser	Gln	Val	Gln	Leu	Gln	Gln	Pro	Gly	Ala	Glu	Leu	Val	Lys	Ala	Gly	Ala	Ser
CTG	TCC	CAG	GTA	CAA	CTG	CAG	CAG	CCT	GGG	GCT	GAG	CTG	GTG	AAG	CCT	GGG	GCC	TCA
	2460			2469				2478			2487			2496			2505	

20		25		30	31	CDR1		35	36									
Val	Lys	Met	Ser	Cys	Lys	Ala	Ser	Gly	Tyr	Thr	Phe	Thr	Ser	Tyr	Asn	Met	His	Trp
GTG	AAG	ATG	TCC	TGC	AAG	GCT	TCT	GGC	TAC	ACA	TTT	ACC	AGT	TAC	AAT	ATG	CAC	TGG
	2517			2526				2536			2544			2553			2562	

40	FR2		45		49	50	52	52A	53	54								
Val	Lys	Gln	Thr	Pro	Gly	Arg	Gly	Leu	Glu	Trp	Ile	Gly	Ala	Ile	Tyr	Pro	Gly	Asn
GTA	AAA	CAG	ACA	CCT	GGT	CGG	GGC	CTG	GAA	TGG	ATT	GGA	GCT	ATT	TAT	CCC	GGA	AAT
	2574			2583				2592			2601			2610			2619	

55	CDR2		60		65	66	FR3		70									
Gly	Asp	Thr	Ser	Tyr	Asn	Gln	Lys	Phe	Lys	Gly	Lys	Ala	Thr	Leu	Thr	Ala	Asp	Lys
GGT	GAT	ACT	TCC	TAC	AAT	CAG	AAG	TTC	AAA	GGC	AAG	GCC	ACA	TTG	ACT	GCA	GAC	AAA
	2631			2640				2649			2658			2667			2676	

75		80		82	82A	82B	82C	83		85								
Ser	Ser	Ser	Thr	Ala	Tyr	Met	Gln	Leu	Ser	Ser	Leu	Thr	Ser	Glu	Asp	Ser	Ala	Val
TCC	TCC	AGC	ACA	GCC	TAC	ATG	CAG	CTC	AGC	AGC	CTG	ACA	TCT	GAG	GAC	TCT	GCG	GTC
	2688			2697				2706			2715			2724			2733	

90		94	95	CDR3		100	100A	100B	100C	100D	101	102	103					
Tyr	Tyr	Cys	Ala	Arg	Ser	Thr	Tyr	Tyr	Gly	Gly	Asp	Trp	Tyr	Phe	Asn	Val	Trp	Gly
TAT	TAC	TGT	GCA	AGA	TCG	ACT	TAC	TAC	GGC	GGT	GAC	TGG	TAC	TTC	AAT	GTC	TGG	GGC
	2745			2754				2763			2772			2781			2790	

105	FR4		110		113			
Ala	Gly	Thr	Thr	Val	Thr	Val	Ser	Ala
GCA	GGG	ACC	ACG	GTC	ACC	GTC	TCT	GCA
	2802			2811				2820

FIG. 5




FIG. 6




FIG. 7






FIG. 8




FIG. 9A




FIG. 9B






FIG. 9C

*FIG. 10**FIG. 11*




FIG. 12




FIG. 13




FIG. 14A




FIG. 14B



European Patent  
Office

# EUROPEAN SEARCH REPORT

Application Number  
EP 96 20 0772

DOCUMENTS CONSIDERED TO BE RELEVANT			
Category	Citation of document with indication, where appropriate, of relevant passages	Relevant to claim	CLASSIFICATION OF THE APPLICATION (Int.Cl.6)
X,D	EP-A-0 274 394 (INTERNATIONAL GENETIC ENGINEERING, INC (INGENE)) * page 9, line 22 - line 37 * * page 13, line 8 - line 44 * ---	1	A61K39/395 A61K51/10 C12N15/02 C12P21/08 C07K16/46 //C07K16/28
X,D	JOURNAL OF IMMUNOLOGY, vol. 139, no. 10, 15 November 1987, BALTIMORE US, pages 3521-3526, XP002010074 ALVIN Y. LIU ET AL.: "PRODUCTION OF A MOUSE-HUMAN CHIMERIC MONOCLONAL ANTIBODY TO CD20 WITH POTENT FC-DEPENDENT BIOLOGIC ACTIVITY." * the whole document * ---	1	
X,P	THE NETHERLANDS CANCER INSTITUTE AMSTERDAM ANNUAL REPORT 1993, 1993, pages 47-48, XP000578064 A. HEKMAN ET AL.: "IMMUNOTHERAPY" * the whole document * ---	1-5	
X,P	JOURNAL OF CELLULAR BIOCHEMISTRY, vol. 17, no. SUPPLEMENT 17E, 29 March 1993 - 25 April 1993, page 260 XP000578063 MITCHELL REFF ET AL.: "DEPLETION OF B CELLS IN VIVO BY A CHIMERIC MOUSE HUMAN MONOCLONAL ANTIBODY TO CD20" * paragraph T103 * -----	1-5,9,17	
<div>TECHNICAL FIELDS SEARCHED (Int.Cl.6)</div> <div>C07K</div>			
The present search report has been drawn up for all claims			
Place of search THE HAGUE		Date of completion of the search 2 August 1996	Examiner Rempp, G
<div>CATEGORY OF CITED DOCUMENTS</div> <div> X : particularly relevant if taken alone  Y : particularly relevant if combined with another document of the same category  A : technological background  O : non-written disclosure  P : intermediate document </div> <div> T : theory or principle underlying the invention  E : earlier patent document, but published on, or after the filing date  D : document cited in the application  L : document cited for other reasons  .....  &amp; : member of the same patent family, corresponding document </div>			

EPO FORM 1503 03.82 (P04C01)